









# JOURNAL OF AGRICULTURAL RESEARCH

VOLUME VI

APRIL 3, 1916—SEPTEMBER 25, 1916

DEPARTMENT OF AGRICULTURE  
WASHINGTON, D. C.

PUBLISHED BY AUTHORITY OF THE SECRETARY  
OF AGRICULTURE, WITH THE COOPERATION  
OF THE ASSOCIATION OF AMERICAN AGRICUL-  
TURAL COLLEGES AND EXPERIMENT STATIONS

### EDITORIAL COMMITTEE

---

#### FOR THE DEPARTMENT

KARL F. KELLERMAN, CHAIRMAN

*Physiologist and Assistant Chief, Bureau  
of Plant Industry*

EDWIN W. ALLEN

*Chief, Office of Experiment Stations*

CHARLES L. MARLATT

*Assistant Chief, Bureau of Entomology*

#### FOR THE ASSOCIATION

RAYMOND PEARL

*Biologist, Maine Agricultural Experiment  
Station*

H. P. ARMSBY

*Director, Institute of Animal Nutrition, The  
Pennsylvania State College*

E. M. FREEMAN

*Botanist, Plant Pathologist, and Assistant  
Dean, Agricultural Experiment Station of  
the University of Minnesota*

---

All correspondence regarding articles from the Department of Agriculture should be addressed to Karl F. Kellerman, Journal of Agricultural Research, Washington, D. C.

All correspondence regarding articles from Experiment Stations should be addressed to Raymond Pearl, Journal of Agricultural Research, Orono, Maine.

## CONTENTS

	Page
Relation of Carbon Bisulphid to Soil Organisms and Plant Growth. E. B. FRED.....	1
Climatic Conditions as Related to <i>Cercospora beticola</i> . VENUS W. POOL and M. B. MCKAY.....	21
Soluble Nonprotein Nitrogen of Soil. R. S. POTTER and R. S. SNYDER.....	61
Oviposition of <i>Megastigmus spermatrophus</i> in the Seed of Douglas Fir. J. M. MILLER.....	65
Citrus Canker. FREDERICK A. WOLF.....	69
Determination of Stearic Acid in Butter Fat. E. B. HOLLAND, J. C. REED, and J. P. BUCKLEY, JR.....	101
Life History and Habits of Two New Nematodes Parasitic on Insects. J. H. MERRILL and A. L. FORD.....	115
Insect Injury to Cotton Seedlings. B. R. COAD and R. W. HOWE.....	129
A Sex-Limited Color in Ayrshire Cattle. EDWARD N. WENT- WORTH.....	141
Watermelon Stem-End Rot. F. C. MEIER.....	149
Effect of Pasteurization on Mold Spores. CHARLES THOM and S. HENRY AYERS.....	153
Effect of Water in the Ration on the Composition of Milk. W. F. TURNER, W. H. SHAW, R. P. NORTON, and P. A. WRIGHT.....	167
Crown Gall Studies, Showing Changes in Plant Structures Due to a Changed Stimulus. ERWIN F. SMITH.....	179
Effect of Certain Species of <i>Fusarium</i> on the Composition of the Potato Tuber. LON A. HAWKINS.....	183
<i>Hyperaspis binotata</i> , a Predatory Enemy of the Terrapin Scale. F. L. SIMANTON.....	197
Test of Three Large-Sized Reinforced-Concrete Slabs under Con- centrated Loading. A. T. GOLDBECK and E. B. SMITH.....	205
Occurrence of Sterile Spikelets in Wheat. A. E. GRANTHAM and FRAZIER GROFF.....	235
Effect of Cold-Storage Temperatures upon the Pupæ of the Medi- terranean Fruit Fly. E. A. BACK and C. E. PEMBERTON.....	251
Effect of Climatic Factors on the Hydrocyanic-Acid Content of Sorghum. J. J. WILLAMAN and R. M. WEST.....	261
Egg and Manner of Oviposition of <i>Lyctus planicollis</i> . THOMAS E. SNYDER.....	273
<i>Hypoderma deformans</i> , an Undescribed Needle Fungus of Western Yellow Pine. JAMES K. WEIR.....	277



	Page
Ornix germinatella, the Unspotted Tentiform Leaf Miner of Apple. L. HASEMAN.....	289
A Western Fieldrot of the Irish Potato Caused by Fusarium radicola. O. A. PRATT.....	297
Comparative Study of the Root Systems and Leaf Areas of Corn and the Sorghums. E. C. MILLER.....	311
Production of Clear and Sterilized Anti-Hog-Cholera Serum. M. DORSET and R. R. HENLEY.....	333
Silver-Scurf of the Irish Potato Caused by Sponylocadium atrovirens. EUGENE S. SCHULTZ.....	339
Woolly Pear Aphis. A. C. BAKER and W. M. DAVIDSON.....	351
Pathological Histology of Strawberries Affected by Species of Botrytis and Rhizopus. NEIL E. STEVENS.....	361
Life Histories and Methods of Rearing Hessian-Fly Parasites. C. M. PACKARD.....	367
Effect of Röntgen Rays on the Tobacco, or Cigarette, Beetle and the Results of Experiments with a New Form of Röntgen Tube. G. A. RUNNER.....	383
Stimulating Influence of Arsenic upon the Nitrogen-Fixing Organisms of the Soil. J. E. GREAVES.....	389
Transmission and Control of Bacterial Wilt of Cucurbits. FREDERICK V. RAND and ELLA M. A. ENLWS.....	417
Correlated Characters in Maize Breeding. G. N. COLLINS.....	435
Comparative Study of the Amount of Food Eaten by Parasitized and Nonparasitized Larvæ of Cirphis unipuncta. DANIEL G. TOWER.....	455
Aleyrodidae, or White Flies Attacking the Orange, with Descriptions of Three New Species of Economic Importance. A. L. QUANTANCE and A. C. BAKER.....	459
Relative Water Requirement of Corn and the Sorghums. EDWIN C. MILLER.....	473
Availability of Mineral Phosphates for Plant Nutrition. W. L. BURLISON.....	485
California Green Lacewing Fly. V. L. WILDERMUTH.....	515
Rape as Material for Silage. A. R. LAMB and JOHN M. EYVARD.....	527
Effect of Autolysis upon Muscle Creatin. RALPH HOAGLAND and C. N. MCBRYDE.....	535
Storage Rots of Economic Aroids. L. L. HARTER.....	549
Experiments with Clean Seed Potatoes on New Land in Southern Idaho. O. A. PRATT.....	573
Digestibility of Very Young Veal. C. F. LANGWORTHY and A. D. HOLMES.....	577
Influence of Calcium and Magnesium Compounds on Plant Growth. F. A. WYATT.....	589

	Page
Larval Characters and Distribution of Two Species of <i>Diatraea</i> . T. E. HOLLOWAY.....	621
The Disease of Potatoes Known as "Leak." LON A. HAWKINS.. Digestibility of Hard Palates of Cattle. C. F. LANGWORTHY and A. D. HOLMES.....	627 641
Some Properties of the Virus of the Mosaic Disease of Tobacco. H. A. ALLARD.....	649
Life Cycles of the Bacteria. F. LÖNNIS and N. R. SMITH.....	675
A Respiration Calorimeter, Partly Automatic, for the Study of Metabolic Activity of Small Magnitude. C. F. LANGWORTHY and R. D. MILNER.....	703
Mottle-Leaf of Citrus Trees in Relation to Soil Conditions. LY- MAN J. BRIGGS, C. A. JENSEN, and J. W. McLANE.....	721
Vegetative Succession Under Irrigation. J. FRANCIS MACBRIDE.. Agricultural Value of Impermeable Seeds. GEORGE T. HAR- RINGTON.....	741 761
Mendelism of Short Ears in Sheep. E. G. RITZMAN.....	797
Life-History Studies of <i>Cirphis Unipuncta</i> , the True Army Worm. JOHN J. DAVIS and A. F. SATTERTHWAIT.....	799
Infection of Timothy by <i>Puccinia graminis</i> . E. C. STAKMAN and F. J. PIEMEISEL.....	813
Control of the Powdery Dryrot of Western Potatoes Caused by <i>Fusarium trichothecioides</i> . O. A. PRATT.....	817
Use of the Moisture Equivalent for the Indirect Determination of the Hygroscopic Coefficient. FREDERICK J. ALWAY and JOUETTE C. RUSSEL.....	833
<i>Thersilochus conotracheli</i> , a Parasite of the Plum Curculio. R. A. CUSHMAN.....	847
Effect on Plant Growth of Sodium Salts in the Soil. FRANK B. HEADLEY, E. W. CURTIS, and C. S. SCOFIELD.....	857
Observations on the Life History of the Army Cutworm, <i>Chori- zagrotis auxiliaris</i> . R. A. COOLEY.....	871
<i>Aphidoletes meridionalis</i> , an Important Dipterous Enemy of Aphids. JOHN J. DAVIS.....	883
Influence of Barnyard Manure and Water upon the Bacterial Activities of the Soil. J. E. GREAVES and E. G. CARTER.....	889
Progressive Oxidation of Cold-Storage Butter. D. C. DYER.....	927
Bacteriological Studies of a Soil Subjected to Different Systems of Cropping for Twenty-five Years. P. L. GAINES and W. M. GIBBS.....	953
Studies on the Physiology of Reproduction in the Domestic Fowl.— XV. Dwarf Eggs. RAYMOND PEARL and MAYNIE R. CURTIS.. $\alpha$ -Crotonic Acid, a Soil Constituent. E. H. WALTERS and LOUIS E. WISE.....	977 1043
Index.....	1047

## ERRATA

- Page 151, line 25, "Sphacropsidae" should read "Sphaerioidaceae."
- Page 150, "Penicillium camemberti, var. rogeri" should read "Penicillium camemberti, var. rogeri."
- Page 296, Pl. XXXIII, figs. 3 to 15. The magnification of the illustrations should be half that stated in the legend.
- Page 303, line 17, "Plate XXXVI, figures 1 to 4" should read "Plate XXXVII, figures 1 to 4" and "In figure 4, Plate XXXVI" should read "In figure 4, Plate XXXVII."
- Page 318, Table IV, under head "General remarks," "rooting" should read "shootings."
- Page 337, Table II, 4th column, "Phenolized defibrinated blood 3895 (unwashed)" should read "Phenolized defibrinated blood 3895."
- Page 377, last line, 2d paragraph, "winged" should read "wingless."
- Page 384, Table I, 6th column, "Current (milliamperes minutes)" should read "Current (milliamperes)."
- Page 388, line 13 from bottom, omit "with humidity at 57."
- Page 419, line 25, "The twelve-spotted (or squash) lady beetle" should read "The squash lady beetle."
- Page 419, line 28, "(*Crepidodera cucumeris*)" should read "(*Epitrix cucumeris*)."
- Page 459, lines 2 and 24, omit "Three."
- Page 471, line 4, "*Aleurodes mori* Ckll." should read "*Aleurodes mori*, var. *arizomenae* Ckll."
- Page 762, Table I, first column, "*Medicago arborea*" should read "*Medicago arabica*."
- Page 791, Table XIV, 1st column, "(p. 21)" should read "(p. 781)."
- Page 866, legend under figure 5, "The solid black line, etc." should read "The hatched line."
- Page 866, legend under figure 6, omit sentences 2 and 3.
- Page 881, line 6 from bottom, "April" should read "May."

# ILLUSTRATIONS

## PLATES

### RELATION OF CARBON BISULPHID TO SOIL ORGANISMS AND PLANT GROWTH

	Page
PLATE I. Plate cultures of soil organisms growing on agar: Fig. 1.—Colonies of organisms from untreated soil. Fig. 2.—Colonies from soil treated with 2 per cent of carbon bisulphid. Fig. 3.—Colonies from soil treated with 2 per cent of carbon bisulphid and evaporated. Fig. 4.—Colonies from soil treated with 2 per cent of carbon bisulphid, evaporated, and reinoculated with 5 per cent of soil from an untreated jar.....	20
PLATE II. Fig. 1.—Effect of varying amounts of carbon bisulphid on mustard; A, B, soil untreated; C, D, soil treated with 0.5 per cent of carbon bisulphid; E, F, soil treated with 1 per cent of carbon bisulphid; G, H, soil treated with 2 per cent of carbon bisulphid. Fig. 2.—Effect of carbon bisulphid on clover in peat soil; A, B, soil untreated; C, D, soil treated with 2 per cent of carbon bisulphid. Fig. 3.—Effect of carbon bisulphid on buckwheat in sand cultures; A, B, soil untreated; C, D, soil treated with 2 per cent of carbon bisulphid.....	20

### CLIMATIC CONDITIONS AS RELATED TO *CERCOSPORA BETICOLA*

PLATE III. <i>Cercospora beticola</i> : Overwintering tests on the experimental field at Rocky Ford, Colo., during 1912-13: Sugar-beet leaves infected with <i>Cercospora beticola</i> (1) stored in soil in boxes, (2) buried in the ground at different depths from 1 to 8 inches, and (3) left exposed above the ground in a pile of hayed sugar-beet tops.....	60
PLATE IV. Field stations for the collection of weather data at Rocky Ford, Colo., in 1913: Fig. 1.—Weather shelter, anemometer, and rain gauge at edge of sugar-beet field. Fig. 2.—Weather shelter among beet plants, showing hygrothermograph and cog psychrometer. Fig. 3.—Weather shelter of the local Weather Bureau station about 3 miles from sugar-beet field.....	60

### OVIPOSITION OF *MEGASTIGMUS SPERMOTROPHUS* IN THE SEED OF DOUGLAS FIR

PLATE V. Oviposition of <i>Megastigmus spermotrophus</i> in the cones of Douglas fir: Fig. 1.—Type of cage in which the oviposition of <i>Megastigmus spermotrophus</i> was observed. Fig. 2. 3.—Female resting on cone with ovipositor inserted.....	68
PLATE VI. Oviposition of <i>Megastigmus spermotrophus</i> in the cones of Douglas fir: Figs. 1, 2.—Two positions of female on surface of cone with ovipositor inserted. Fig. 3.—Female resting on cone with ovipositor inserted.....	68
PLATE VII. Oviposition of <i>Megastigmus spermotrophus</i> in the cones of Douglas fir: Fig. 1.—Female in act of withdrawing ovipositor from cone. Fig. 2.—Section through a Douglas fir cone on which a female has been killed while in the act of ovipositing. Fig. 3.—A portion of same cone and dead female with ovipositor inserted.....	68

## CITRUS CANKER

	Page
PLATE VIII. Fig. 1.—Grapefruit leaf showing young Citrus cankers. Fig. 2.—Old Citrus canker on Satsuma leaves. Figs. 3, 4.—Seedling grapefruit branches affected with Citrus canker. Fig. 5.—Severe canker infection of branches of <i>Citrus trifoliata</i> .....	100
PLATE IX. Fig. 1.—View of lower side of leaves of seedling grapefruit artificially inoculated with <i>Pseudomonas citri</i> . Fig. 2.—Top view of plant shown in figure 1. Fig. 3.—Spongy white cankers on leaf and twig of seedling grapefruit produced by artificial inoculation. Fig. 4.—Citrus canker on Satsuma leaves resulting from artificial inoculation with <i>Pseudomonas citri</i> . Fig. 5.—Photomicrograph of section of young, open canker on grapefruit.....	100
PLATE X. Fig. 1.—Natural Citrus canker infection on leaves of <i>Citrus trifoliata</i> . Fig. 2.—Mature cankers on fruit of <i>Citrus decumana</i> . Fig. 3.—Canker on seedling grapefruit leaves, entrance having been effected through abrasions made by thorns. Fig. 4.—Young spongy cankers on fruit of <i>Citrus decumana</i> . Fig. 5.— <i>Phoma socia</i> on cellulose agar showing dissolution of cellulose. Fig. 6.—Mature cankerous areas on leaves of Duncan grapefruit....	100
PLATE XI. Fig. 1.—Cankers on old grapefruit leaves which have enlarged during the second growing season. Fig. 2.—Citrus canker resulting from immersion of leaves in a bacterial suspension. Fig. 3.—Culture of <i>Phoma socia</i> showing pycnidial formation in concentric rings. Fig. 4.—Dilution poured plate of <i>Pseudomonas citri</i> on green-bean agar. Fig. 5.—Photomicrograph of pycnidium of <i>Phoma socia</i> taken in reflected sunlight. Fig. 6. Photomicrograph of pycnidia of <i>Phoma socia</i> taken in diffuse light.....	100

## INSECT INJURY TO COTTON SEEDLINGS

PLATE XII. Fig. 1.—Cutworm injury to cotton seedlings; produced in breeding cages. Figs. 2, 3.—Cutworm injury to cotton seedling.....	140
PLATE XIII. Fig. 1.—Cutworm injury to cotton seedling. Fig. 2.—Tussock larva feeding upon cotton leaf. Fig. 3.—Injury produced by a nearly full-grown tussock larva when confined in a screen cage containing potted cotton plants.....	140
PLATE XIV. Cotton leaves showing grasshopper injury.....	140
PLATE XV. Fig. 1.—Underside of cotton leaf showing grasshopper injury. Fig. 2.—Cotton leaf showing grasshopper injury.....	140
PLATE XVI. Fig. 1.—Injury to terminal bud of cotton by lepidopterous larva. Fig. 2.—Two cotton plants from laboratory garden with leaves removed..	140

## WATERMELON STEM-END ROT

PLATE XVII. Watermelons, showing the effect of inoculation with species of <i>Diplodia</i> : Fig. 1.—The upper four melons were held as checks; the lower five are melons nine days after having been inoculated with a culture of <i>Diplodia</i> sp. Fig. 2.—A watermelon nine days after having been inoculated with a culture of <i>Diplodia tubercicola</i> E. and E.....	152
--	-----

## CROWNCALL STUDIES SHOWING CHANGES IN PLANT STRUCTURES DUE TO A CHANGED STIMULUS

PLATE XVIII. Teratoid crown-galls produced in <i>Pelargonium</i> spp.....	182
PLATE XIX. Teratoid crown-galls produced in castor-oil plant. Fig. A.—A red-stem variety. Leaves reflexed; axis distorted; and feeble shoots developing out of the axillary tumors. Fig. B.—A green-stem glaucous variety.....	182

	Page
PLATE XX. Teratoid crown-galls produced in tobacco by inoculating <i>Bacterium tumefaciens</i> .....	182
PLATE XXI. The teratoid tumor strand of Plate XX, which gives rise during its course to more than 30 small tumors. Top.—Cross section of outer part of right side of stem of tobacco plant shown on Plate XX. P, outer edge of the phloem; E, epidermis; T St, tumor strand, which is bedded in the normal cortex of the stem. Bottom.—Longitudinal section from upper part of the above tumor strand. The pathological tissues are S T, sieve tubes; C, cambium; Tr, trachei; Sp, spiral vessels.....	182
PLATE XXII. Teratoid crown-galls produced in a tobacco plant by inoculating <i>Bacterium tumefaciens</i> into the leaf axils.....	182
PLATE XXIII. Teratoid crown-galls produced in tobacco leaves with the hop strain of <i>Bacterium tumefaciens</i> by local inoculations—that is, inoculation in places where shoot anlage are not known to exist. Fig. A.—Portion of an upper leaf showing four shoot-bearing tumors growing from upper surface of the inoculated midrib. Fig. B.—Same as A, but the leaf reversed and the midrib stripped of its blade to show two other shoot-bearing tumors which have developed from its under surface. Fig. C.—From middle of another leaf on the same plant as A, but further magnified and photo made on an orthochromatic plate to show the pale green character of the shoot as contrasted with the dark green of the surrounding leaf.....	182
HYPERASPIS BINOTATA, A PREDATORY ENEMY OF THE TERRAPIN SCALE	
PLATE XXIV. <i>Hyperaspis binotata</i> : Fig. 1.—Male, showing the characteristic markings. Fig. 2.—Female, showing the dorsal view. Fig. 3.—Eggs and a second-instar larva. a, Second-instar larva as disclosed by displacing the host; b, larvæ of the terrapin scale, <i>Eulecanium nigrofasciatum</i> ; c, a displaced scale; d, eggs "in situ"; e, egg. Fig. 4.—First-instar larva. Fig. 5.—Method of attacking the mature scales during the third and fourth instars.....	204
PLATE XXV. <i>Hyperaspis binotata</i> : Fig. 1.—Mature larva as it appears when attacking the leaf-attached larvæ of the terrapin scale, <i>Eulecanium nigrofasciatum</i> . Fig. 2.—Ventral view of mature larva. Fig. 3.—Dorsal view of pupa, showing the last larval molt skin and the depressed segmented area. Fig. 4.—Ventral view of pupa.....	204
TESTS OF THREE LARGE-SIZED REINFORCED-CONCRETE SLABS UNDER CONCENTRATED LOADING	
PLATE XXVI. Fig. 1.—Load-applying and load-measuring apparatus for testing reinforced-concrete slabs, showing set-up for 4-point loading. Fig. 2.—Load-measuring apparatus and hydraulic jack for testing reinforced-concrete slabs.....	234
OCCURRENCE OF STERILE SPIKELETS IN WHEAT	
PLATE XXVII.—Comparison of the number of sterile spikelets on bearded and beardless varieties of wheat.....	250
EGG AND MANNER OF OVIPOSITION OF LYCTUS PLANICOLLIS	
PLATE XXVIII. <i>Lyctus planicollis</i> : Fig. 1.—Outline of the egg, showing strand attachment. Fig. 2.—Greatly enlarged view of end of egg, showing granular appearance. Fig. 3.—Larva within egg, ready to hatch. Fig. 4.—Sketch of egg in pore of wood on radial section of green-ash ladder-rung stock, showing longitudinal striæ.....	276

	Page
PLATE XXIX. <i>Lyctus planicollis</i> : Larval burrows in an ash shovel handle. . . . .	276
PLATE XXX. <i>Lyctus planicollis</i> : Pupal cells in "powder-posted" white-ash shovel handle. . . . .	276
PLATE XXXI. <i>Lyctus planicollis</i> : Exit holes of adults in ash shovel handles. . . . .	276

#### HYPODERMA DEFORMANS, AN UNDESCRIBED NEEDLE FUNGUS OF THE WESTERN YELLOW PINE

PLATE XXXII. Fig. 1.—Needles of <i>Pinus ponderosa</i> infected with <i>Hypoderma deformans</i> , showing the apothecia. Fig. 2.—Branches of <i>Pinus ponderosa</i> deformed and broomed by <i>Hypoderma deformans</i> . Fig. 3.—A branch of <i>Pinus ponderosa</i> , showing how it will hang vertically when supporting a large broom caused by <i>Hypoderma deformans</i> . . . . .	288
--	-----

#### ORNIX GEMINATELLA, THE UNSPOTTED TENTIFORM LEAF MINER OF APPLE

PLATE XXXIII. <i>Ornix geminatella</i> Pack.: Fig. 1.—Moth expanded. Fig. 2.—Moth at rest on leaf. Fig. 3.—Egg on lower surface of leaf; also tunnel made by miner on leaving the egg. Fig. 4.—Dorsal view of first larval stage; below, side view of head and thorax. Fig. 5.—Dorsal view of second larval stage. Fig. 6.—Side view of second larval stage. Fig. 7.—Dorsal view of third larval stage, showing edge of thoracic legs. Fig. 8.—Dorsal view of fourth larval stage. Fig. 9.—Side view of fourth larval stage. Fig. 10.—Ventral view of pupa. Fig. 11.—Dorsal view of same. Fig. 12.—Lower surface of leaf with numerous partly developed mines; also two cocoons, one exposed. Fig. 13.—Portion of leaf showing a mine in process of development. Fig. 14.—A small twig showing leaves badly curled and injured by numerous mines. Fig. 15.—Leaf much distorted with 10 mines almost completed; also one cocoon appears at the tip of the leaf. . . . .	296
--	-----

#### A WESTERN FIELDROT OF THE IRISH POTATO TUBER CAUSED BY *FUSARIUM RADICICOLA*

PLATE XXXIV. Fig. 1, 2, 3, 4.—Types of stem-end blackrot, lenticel rot, and eye rot in Idaho Rural potato tubers. Fig. 5, 6.—Longitudinal and cross sections of an Idaho Rural tuber infected with blackrot. . . . .	310
PLATE XXXV. Fig. 1.—Netted Gem potato tuber infected with jelly-end rot. Fig. 2.—Stem-end view of a Netted Gem tuber infected with jelly-end rot. Fig. 3.—Longitudinal section of a Netted Gem tuber infected with jelly-end rot. Fig. 4.—Idaho Rural tuber infected with stem-end and lenticel blackrot, after having been kept 10 days in a moist chamber. Fig. 5.—Idaho Rural tuber infected with lenticel blackrot after having been kept in a moist chamber for 10 days. . . . .	310
PLATE XXXVI. Fig. 1, 2.—Stem-end blackrot produced by stem-end punctures with a needle carrying <i>Fusarium radicicola</i> . Netted Gem and Idaho Rural potato tubers. Fig. 3.—Lenticel blackrot produced by spraying the tuber with a spore suspension of <i>F. radicicola</i> . Netted Gem tuber. Fig. 4.—Same tuber as shown in figure 3; after having been kept a few days longer in the moist chamber. Fig. 5.—Stem-end blackrot produced by spraying an Idaho Rural tuber with a spore suspension of <i>F. radicicola</i> . Fig. 6.—Stem-end blackrot produced by the inoculation of the tuber stolon. Idaho Rural tuber. Fig. 7.—Blackened vascular system produced by the inoculation of the tuber stolon. Idaho Rural tuber. . . . .	310

	Page
PLATE XXXVII. Fig. 1, 2, 3.—Stem-end and lenticel blackrot produced by spraying the growing tubers with a spore suspension of <i>Fusarium radiclecola</i> . Idaho Rural potato tubers. Fig. 4.—Eye infection produced by spraying the growing tuber with a spore suspension of <i>F. radiclecola</i> . Netted Gem tuber. Fig. 5, 6, 7.—Stem-end blackrot produced by the inoculation of the stolons of growing Idaho Rural tubers. Fig. 8.—Stem-end rot of Netted Gem tuber produced by inoculating the stolon of the growing tuber. . . . .	310
COMPARATIVE STUDY OF THE ROOT SYSTEMS AND LEAF AREAS OF CORN AND THE SORGHUMS	
PLATE XXXVIII. Fig. 1.—Method used in isolating root systems in the field. View of two soil prisms ready for washing. Fig. 2.—Method used in isolating root systems. Fig. 3.—Method of washing used in the isolation of the root systems. . . . .	332
PLATE XXXIX. Fig. 1.—Root system of a corn plant that had reached a height of 3 feet 6 inches. Fig. 2.—Root systems of two corn plants with a height of 1 foot 6 inches. Fig. 3.—Root system of a Dwarf milo plant at the age of 4 weeks. Fig. 4.—Root systems of two Blackhull kafir plants 1 foot in height. . . . .	332
PLATE XL. Fig. 1.—Root systems of two mature corn plants. Fig. 2.—Root system of a corn plant at the time of "shooting". . . . .	332
PLATE XLI. Fig. 1.—Root systems of two Blackhull kafir plants at the time they had reached a height of 6 feet and were blooming. Fig. 2.—Root system of two Dwarf milo plants at the time the seed was in the milk stage. . . . .	332
PLATE XLII. Fig. 1.—Portion of a primary root of Pride of Saline corn, showing the number and relative size of the secondary roots. Fig. 2.—Portions of the primary roots of Blackhull kafir, showing the number and relative size of the secondary roots. . . . .	332
PLATE XLIII. Fig. 1.—Pride of Saline corn, Dwarf milo, and Blackhull kafir plants, showing their relative leaf and sheath areas at 4 weeks of age. Fig. 2.—Pride of Saline corn, Dwarf milo, and Blackhull kafir plants, showing their relative leaf and sheath areas at 6 weeks of age. . . . .	332
PLATE XLIV. Fig. 1.—Pride of Saline corn, Dwarf milo, and Blackhull kafir plants, showing their relative leaf and sheath areas at 8 weeks of age. Fig. 2.—Pride of Saline corn, Dwarf milo, and Blackhull kafir plants, showing their relative leaf and sheath areas at 10 weeks of age. . . . .	332
SILVER-SCURF OF THE IRISH POTATO CAUSED BY <i>SPONDYLOCLADIUM ATROVIRENS</i>	
PLATE XLV. Fig. 1.—Potato tubers showing shriveling and a silvery appearance caused by <i>Spondylocladium atrovirens</i> . Fig. 2.—Tuber naturally infected by <i>S. atrovirens</i> , showing the segregated area type of infection. Fig. 3.—Immature potato tuber artificially inoculated with conidia of <i>S. atrovirens</i> , July, 1913, at Houlton, Me. . . . .	350
PLATE XLVI. Fig. 1.—Photomicrograph of <i>Spondylocladium atrovirens</i> on corn-meal agar, showing method of development of conidiophores and conidia in the early stages. Fig. 2.—Photomicrograph of <i>S. atrovirens</i> in hanging-drop culture, showing development of conidiophore and conidia in mature stages. Fig. 3.—Negative heliotropism of <i>S. atrovirens</i> on corn-meal agar exposed on one side to daylight from April 8 to April 24, 1915, in laboratory at room temperature. . . . .	350
PLATE XLVII. Effect of temperature upon mycelial development of <i>Spondylocladium atrovirens</i> in pure culture on corn-meal agar at end of four weeks. . . . .	350



	Page
PLATE XLVIII. Contact infection. A part of the new tubers becoming infected with <i>Spondylocladium atrovirens</i> by means of contact with the infected mother tuber.....	350
PATHOLOGICAL HISTOLOGY OF STRAWBERRIES AFFECTED BY SPECIES OF BOTRYTIS AND RHIZOPUS	
PLATE XLIX. A-H, Strawberry cells attacked by <i>Botrytis</i> sp.: A, Hypha growing partly between and partly within strawberry cells; B, hyphae inside strawberry cells in which remnants of the protoplasm may still be distinguished; C, hypha passing from one cell into another across a short intercellular space; D, E, F, G, H, hyphae entering cells in various ways (in G the hypha has pushed back a portion of the cell wall before breaking through). I-M, Strawberry cells attacked by <i>Rhizopus</i> sp.: I, Hypha growing between the epidermis and the adjacent layer of storage cells; J, hypha growing over the surface of the strawberry; K, hyphae growing underneath the epidermal layer and between the storage cells; L, <i>Rhizopus</i> sp. growing between epidermal cells; M, germinating spore in cavity formed by a seed.....	366
PLATE L. Strawberry cells attacked by <i>Rhizopus</i> sp. A, Normal storage cell of strawberry; B, storage cell showing a slight contraction of the protoplasm; C, D, E, F, G, progressive contraction of protoplasm of host cells near hyphae; H, I, J, strawberry cells near hyphae in which the cell wall has crumpled with the contraction of the protoplasm; K, M, hyphae inside cells; L, hyphae growing between cells of the strawberry; K, L, M are drawn from berries which had been rotted in the desiccator.....	366
LIFE HISTORIES AND METHODS OF REARING HESSIAN-FLY PARASITES	
PLATE LI. Fig. 1.—Egg of <i>Eupelmus allynii</i> . Fig. 2.—Egg of <i>Eupelmus allynii</i> in situ. Figs. 3, 4.—Pupa of <i>Eupelmus allynii</i> . Fig. 5.—Egg of <i>Merisus destructor</i> . Fig. 6.—Pupa of <i>Merisus destructor</i> . Fig. 7.—Egg of <i>Micromelus subapterus</i> . Fig. 8.—Pupa of <i>Micromelus subapterus</i> .....	382
PLATE LII. Fig. 1.—Mandibles of full-grown larva of <i>Eupelmus allynii</i> . Fig. 2.—Larva of <i>Eupelmus allynii</i> . Fig. 3.—Mandibles of full-grown larva of <i>Merisus destructor</i> . Fig. 4.—Larva of <i>Merisus destructor</i> . Fig. 5.—Mandibles of full-grown larva of <i>Micromelus subapterus</i> . Fig. 6.—Larva of <i>Micromelus subapterus</i> .....	382
TRANSMISSION AND CONTROL OF BACTERIAL WILT OF CUCURBITS	
PLATE LIII.—Two wilted cucumber plants which contracted bacterial wilt at beetle gnawings of the leaves marked x. Three healthy, uninjured plants in same hill are also shown.....	434
PLATE LIV.—Plots in field 1, East Marion, Long Island, N. Y., 1915. Fig. 1.—Plot sprayed with Bordeaux mixture and lead arsenate, beginning June 25. Fig. 2.—Plot sprayed with Bordeaux mixture and lead arsenate, beginning July 19, after most of the striped-beetle injury had occurred. Fig. 3.—Plot sprayed with Bordeaux mixture and lead arsenate, beginning July 27. Fig. 4.—General view of field, showing cages and meteorological-instrument shelter.....	434
CORRELATED CHARACTERS IN MAIZE BREEDING	
PLATE LV.—Typical plant of the Waxy Chinese variety of maize, showing numerous tassel branches, erect leaf blades, one-sidedness, and curved tassel.....	454

	Page
PLATE LVI. Fig. 1.—Uppermost leaf sheaths of Chinese maize plant, showing the one-sided arrangement of leaf blades and absence of hairs. Fig. 2.—Leaf sheath of the Waxy Chinese variety of maize, showing the transverse lines and absence of hairs. ....	454
PLATE LVII.—A plant of the Esperanza variety of maize, showing the drooping leaves, few tassel branches, and elongated internodes characteristic of the variety. ....	454
PLATE LVIII.—Leaf sheaths of the Esperanza variety of maize, showing the maximum development of tuberculate hairs. ....	454
PLATE LIX.—A leaf sheath of a second-generation hybrid maize plant. ....	454
PLATE LX.—A first-generation plant of Chinese×Esperanza maize hybrid. ....	454
PLATE LXI.—A second-generation plant of a Chinese×Esperanza maize hybrid. ....	454
PLATE LXII.—A second-generation plant of a Chinese×Esperanza maize hybrid. ....	454
PLATE LXIII.—A second-generation plant of a maize hybrid, showing the "goose-neck" character that appeared for the first time in this hybrid. ....	454

ALCYRODIDAE, OR WHITE FLIES ATTACKING THE ORANGE, WITH DESCRIPTIONS OF THREE NEW SPECIES OF ECONOMIC IMPORTANCE

PLATE LXIV.— <i>Aleurocanthus woglumi</i> : Eggs, larvæ, and pupa cases on orange leaves. ....	472
PLATE LXV.— <i>Aleurocanthus woglumi</i> : Fig. 1.—Colony on an orange leaf. Fig. 2.—Eggs and pupa cases. ....	472
PLATE LXVI.—Fig. 1.— <i>Dialeurodes citri</i> : Pupæ. Fig. 2.—Male and female adults of an alcyrodid. Fig. 3.— <i>Aleurolobus marlattii</i> . ....	472
PLATE LXVII. <i>Aleurothrixus howardi</i> : Larvæ and pupa cases on an orange leaf. ....	472
PLATE LXVIII. <i>Aleurothrixus porteri</i> : Larvæ and pupa cases on <i>Myrtus</i> sp. ....	472
PLATE LXIX. Fig. 1.— <i>Tetraleurodes mori</i> , var. <i>arizonensis</i> : Larvæ and pupa cases on an orange leaf. Fig. 2.— <i>Tetraleurodes mori</i> : Pupa cases on a mulberry leaf. ....	472

RELATIVE WATER REQUIREMENT OF CORN AND THE SORGHUMS

PLATE LXX. Fig. 1.—General view of the screened inclosure and the scale house. Fig. 2.—Method of moving the cans. Fig. 3.—General view of the plant shelter and the surrounding country at Garden City, Kans. ....	484
PLATE LXXI. Fig. 1.—Dwarf milo, grown May 22 to September 3, 1915. Fig. 2.—Dwarf Blackhull kafir, grown May 22 to September 11, 1915. Fig. 3.—Feterita, grown May 22 to September 6, 1915. ....	484
PLATE LXXII. Fig. 1.—Sudan grass, grown May 22 to September 14, 1915. Fig. 2.—Pride of Saline corn, grown May 22 to August 25, 1915. Fig. 3.—Blackhull kafir, grown May 22 to September 18, 1915. Fig. 4.—Method of sealing the lids with tape and the wax seal around the plants. ....	484

AVAILABILITY OF MINERAL PHOSPHATES FOR PLANT NUTRITION

PLATE LXXIII. Effect of varying quantities of Tennessee brown rock phosphates on plant growth: Fig. 1.—Spring wheat. Fig. 2.—Sixty-Day oats. ....	514
PLATE LXXIV. Effect of varying quantities of Tennessee brown rock phosphate on plant growth: Fig. 1.—Barley. Fig. 2.—Timothy. ....	514
PLATE LXXV. Effect of varying quantities of Tennessee brown rock phosphate on plant growth: Fig. 1.—Cowpeas. Fig. 2.—Soybeans. Fig. 3.—Red clover. Fig. 4.—Alfalfa. ....	514
PLATE LXXVI. Effect of different kinds of mineral phosphate applied in different quantities for red clover. ....	514

	Page
PLATE LXXVII. Cowpeas, showing the comparative effect of Tennessee brown rock phosphate alone and in combination with dextrose.....	514
PLATE LXXVIII. Cowpeas, showing the comparison of their growth when treated with Tennessee brown rock phosphate, phosphate and dextrose, and phosphate, dextrose, and calcium carbonate.....	514
PLATE LXXIX. Effect of different substances on the growth of cowpeas: Fig. 1.—Growth after the addition of varying quantities of raw rock. Fig. 2.—Growth after the addition of dextrose and soluble phosphate.....	514
PLATE LXXX. Effect of various substances and combinations on the growth of cowpeas: Fig. 1.—Effect of adding lime, phosphate rock, dextrose and lime, and phosphate rock, dextrose, and lime to the soil. Fig. 2.—Effect of adding nothing, lime, phosphate rock, and phosphate rock and lime to the soil.....	514
STORAGE-ROTS OF ECONOMIC AROIDS.	
PLATE LXXXI. Fig. 1.—A dasheen corm ( <i>Colocasia esculenta</i> ) showing Java blackrot produced by <i>Diplodia tubericola</i> . Fig. 2.—A corm of <i>Alocasia sp.</i> showing Java blackrot produced by <i>D. tubericola</i> . Fig. 3.—A dasheen tuber partially decayed by <i>Sclerotium rolfsii</i> .....	572
PLATE LXXXII. Fig. 1.—A tuber of <i>Colocasia esculenta</i> showing a powdery grayrot caused by <i>Fusarium solani</i> . Fig. 2.—A tuber of <i>Xanthosoma sagittifolium</i> showing partial decay by <i>Fusarium solani</i> . Fig. 3.—A tuber of <i>C. esculenta</i> softened throughout by <i>Sclerotium rolfsii</i> .....	572
PLATE LXXXIII. A corm of <i>Colocasia esculenta</i> from Brooksville, Fla., mostly rotted away by <i>Bacillus carotovorus</i> .....	572
INFLUENCE OF CALCIUM AND MAGNESIUM COMPOUNDS ON PLANT GROWTH	
PLATE LXXXIV. Fig. 1.—Growth of wheat in sand containing varying quantities of calcium and magnesium. Fig. 2.—Growth of alfalfa in sand containing varying amounts of calcium and magnesium.....	620
PLATE LXXXV. Fig. 1.—Growth of soybeans following a crop which had already absorbed most of the readily available calcium and magnesium. Fig. 2.—Growth of soybeans in soil treated with magnesium.....	620
PLATE LXXXVI. Fig. 1.—Comparative growth of soybeans in brown silt loam and dolomite, showing that the loam would have been improved by the addition of some limestone or dolomite. Fig. 2.—Soybeans in sand treated with magnesium, showing that their growth increases inversely with the quantity of magnesium applied as sulphate.....	620
PLATE LXXXVII. Comparative root production of wheat grown in the chlorides, sulphates, and carbonates of magnesium and calcium.....	620
PLATE LXXXVIII. Fig. 1.—Comparative growth of wheat in sand and in dolomite. Fig. 2.—Comparative growth of wheat in magnesium chloride and magnesium sulphate.....	620
LARVAL CHARACTERS AND DISTRIBUTION OF TWO SPECIES OF DIATRAEA	
PLATE LXXXIX. Fig. 1.— <i>Diatraea saccharalis crambidoides</i> : Larva, summer form, dorsal view. Fig. 2.— <i>D. zeacolella</i> : Larva, summer form, dorsal view. Fig. 3.— <i>D. saccharalis crambidoides</i> : Larva, summer form, side view. Fig. 4.— <i>D. zeacolella</i> : Larva, summer form, side view. Fig. 5.— <i>D. saccharalis crambidoides</i> : Larva, winter form, dorsal view. Fig. 6.— <i>D. zeacolella</i> : Larva, winter form, dorsal view.....	626

## THE DISEASE OF POTATOES KNOWN AS "LEAK"

PLATE XC. Potatoes affected with potato leak: Fig. 1, 2.—Natural infection from fork wound. Fig. 3.—Rot produced by inoculation with *Pythium debaryumum*. Fig. 4.—Rot produced by inoculation with *Rhizopus nigricans*. Page 640

SOME PROPERTIES OF THE VIRUS OF THE MOSAIC DISEASE OF TOBACCO

PLATE XCI. Livingstone atmometer porous cup as used for filtration Page 674

## LIFE CYCLES OF THE BACTERIA

PLATE A. Fig. 1.—Azotobacter 11. Mannite-nitrate solution, 5 days old. Types A and La. Some cells in conjunction. Fig. 2.—Azotobacter 21. Contact prepare from a colony on mannite agar, 4 days old. Types A, L. Most cells in conjunction. Fig. 3.—Azotobacter 23. Contact prepare from a colony on mannite-agar, 4 days old. Types A, B, I, K $\alpha$ , and many conjunct cells. Fig. 4.—Azotobacter 13. Jannite-nitrate solution, 17 days old. Type K $\lambda$ . Fig. 5.—Azotobacter 14. Mannite-nitrate solution, 5 days old. Type B forming I. Fig. 6.—Azotobacter 8. Beef bouillon. Type B forming types I and J. Page 702

PLATE B. Fig. 7.—Azotobacter 21. Mannite-agar colony, 4 days old. Type C forming types D and I. Fig. 8.—Azotobacter 22. Mannite-agar colony, 4 days old. Type C forming D, also A in conjunction. Fig. 9.—Azotobacter 11. From a filter paper strip in mannite-peptone solution, 16 days old. Types A and B forming D. Fig. 10.—Azotobacter 3. Mannite-peptone solution, 24 days old. Types L and M forming D. Fig. 11.—Azotobacter 11. Mannite-peptone solution, 16 days old. Type D (stained) resulting from Type C. Fig. 12.—Azotobacter 6. From condensation water of mannite-agar slant, 1 day old. Type D (unstained) containing regenerative units. Page 702

PLATE C. Fig. 13.—Azotobacter 24. Mannite-nitrate solution kept 5 days after having been heated 1 minute at 96° C. Types I and F developing from D. Some I germinating in conjunct stage and inclining to form spores. Fig. 14.—Azotobacter 1. Mannite-nitrate solution, 10 days old. Types B, K $\beta$ , E, and Fa developing from stained and unstained type D. Fig. 15.—Azotobacter 15. From condensation water of a mannite-nitrate agar slant, 2 days old. Types Fa and F $\beta$  developing from type D. Fig. 16.—Azotobacter 17. Mannite-soil-extract agar, 2 months old. Type E, Fa, K $\phi$ , and G developing from type D. Fig. 17.—Azotobacter 17. Mannite-nitrate agar, 10 days old. Prepare treated with hot aqueous fuchsin. Type G, partially dissolved; also K $\phi$ . Fig. 18.—Azotobacter 7. Mannite-soil-extract solution, 14 days old. Type H forming D. Page 702

PLATE D. Fig. 19.—Azotobacter 2. Mannite-nitrate agar, 23 days old. Spores forming type D. Fig. 20.—Azotobacter 2. Mannite nitrate-agar, 6 days old. Types L and F, endospores and exospores and dissolving of spores to type D. Fig. 21.—Azotobacter 18. From a filter paper strip in mannite solution, 25, days old. Type L with gonidia, forming B (type J $\lambda$ ). Fig. 22.—Azotobacter 7. Mannite-soil-extract agar, 2 months old. Types E and F forming B. Fig. 23.—Azotobacter 7. Mannite-soil-extract agar, 2 months old. Type B, formed by types E and F, germinating to type G. Fig. 24.—Azotobacter 7. Mannite-soil-extract agar, 2 months old. Type K $\gamma$ . Page 702

	Page
PLATE E. Fig. 25.— <i>Bacillus subtilis</i> (No. 31). Beef agar, 2 days old. Types I and D formed by spores. Fig. 26.— <i>Bacillus subtilis</i> (No. 31). Beef agar, 6 days old. Formation of type I. Fig. 27.— <i>Bacillus subtilis</i> (No. 31). Beef agar, 8 days old. Type I forming H and stained D. Spores forming unstained type D. Fig. 28.—Yellow bacillus (No. 41). Peptone-glycerin solution, 2 days old. Type I germinating from D, stretching to type I. Fig. 29.— <i>Bacterium bulgaricum</i> (No. 49). Whey-yeast agar, 6 days old at 40° C. Types C, D, E, F, G, I, and K. Fig. 30.— <i>Bacterium fluorescens</i> (No. 40). Ammonium-citrate-glycerin solution, 11 days old. Types D and H. ....	702
PLATE F. Fig. 31.— <i>Sarcina flava</i> (No. 43). Beef agar, 1 day old. Type I in conjunction and forming D. Fig. 32.— <i>Streptococcus lactis</i> (No. 48). Peptone lactose solution, 5 days old. Type D, with regenerative units, forming type I. Fig. 33.— <i>Streptococcus lactis</i> (No. 48). Milk, 3 days old. Types D and I in cascin. Fig. 34.— <i>Bacillus radicola</i> (No. 39). Types D and I. Prepare made from a root nodule in 1908. Fig. 35.— <i>Spirillum</i> sp. from Great Salt Lake (No. 46). Beef broth plus 3 per cent of sodium chlorid, 14 days old. Budding and branching forms; stained and unstained regenerative bodies. Some cells in conjunction. Fig. 36.— <i>Spirillum</i> sp. from Great Salt Lake (No. 46). Beef broth plus 3 per cent of sodium chlorid, 14 days old. Type I germinating. ....	702
PLATE G. Fig. 37.— <i>Micrococcus candidans</i> from soil (No. 45). Ammonium-citrate-glycerin solution, 6 days old. Irregular, thick-walled type I. Fig. 38.— <i>Micrococcus candidans</i> from milk (No. 44). Ammonium-citrate-glycerin solution, 2 days old. Irregular, thick-walled type I. Fig. 39.—Yellow bacillus (No. 41). Beef agar, 1 day old. Budding gonidia, formation and germination of type I. Fig. 40.— <i>Bacterium fluorescens</i> (No. 40). Ammonium-citrate-glycerin solution, 2 days old. Budding gonidia, formation and germination of type I. Fig. 41.— <i>Bacterium fluorescens</i> (No. 40). Beef agar, 4 days old. Filterable gonidia germinating. Fig. 42.— <i>Bacterium fluorescens</i> (No. 40). Beef agar, 4 days old. Types D and F formed by filterable gonidia. Dark field. ....	702

A RESPIRATION CALORIMETER, PARTLY AUTOMATIC, FOR THE STUDY OF METABOLIC ACTIVITY OF SMALL MAGNITUDE

PLATE XCII.—General view of the respiration calorimeter: A, Chamber inclosed in heat-insulating cover. B, Tension equalizer to maintain atmospheric pressure in the air of the chamber. C, Absorber table. D, Rotary pump to maintain air circulation. E, Motor to drive pump. F, Bottles containing sulphuric acid to remove water vapor from circulating air. G, Large U-tube, containing soda-lime to remove carbon dioxide from the air. H, Bottle containing sulphuric acid to catch the water vapor from the soda-lime. I, Bottle containing cotton to catch sulphuric acid vapor. J, Small absorbers for determining water vapor and carbon dioxide in residual air. K, Meter to measure the sample of residual air. L, Reservoir to maintain a constant pressure of water in the heat absorber in the chamber. M, Tank to catch water flowing from the heat absorber. N, Pump to raise water from the tank to the reservoir. O, Devices for automatically controlling and recording temperatures. ....

	Page
PLATE XCIII.—Chamber with part of outer covering removed: <i>A</i> , Double metal wall chamber. <i>B</i> , Heat-insulating outer cover. <i>C</i> , Window to chamber. <i>D</i> , Outlet providing passage for pipes, wires, etc., through the walls of the chamber. The exterior ends of the resistance thermometers for ingoing and outgoing water are seen projecting from the outlet. <i>E</i> , Removable top of chamber. <i>F</i> , Device for heating the air entering the respiration chamber. <i>G</i> , Small pipe carrying water for cooling the outer metal wall of the chamber. <i>H</i> , Electric-resistance wire carrying current for heating the outer wall.....	720
PLATE XCIV.—Apparatus connected with the respiration calorimeter: <i>A</i> , Tension equalizer. <i>B</i> , Mixing bottle for equalizing the temperature of water entering the heat absorber. <i>C</i> , Device for heating air entering the respiration chamber. <i>D</i> , Preheater, and <i>E</i> , final heater, for raising the temperature of water entering the heat absorbers. There is an electric-heating coil in the lower half and an electric-resistance thermometer in the upper half of the final heater. <i>F</i> , Temperature indicator comprising part of the apparatus for controlling the temperature of the water entering the heat absorber. <i>G</i> , Multiple-point switch for connecting the resistance thermometers for the metal walls and air of the chamber with the Wheatstone bridge for measuring their temperatures. <i>H</i> , Tube conducting air from the respiration chamber to the rotary air pump. <i>I</i> , Tube conducting air from the purifying devices to the respiration chamber.....	720
PLATE XCV.—Devices for controlling and recording temperatures: <i>A</i> , Mechanism for shifting the contact on the rheostats controlling the current for heating the outer walls of the calorimeter chamber and the ingoing air. <i>B</i> , Ratio coils for the four bridges governing the action of the shifting mechanism <i>A</i> are combined in this box, together with means for checking the constancy of the resistance of the coils and for correcting slight inequalities in them and also to compensate for small differences in the pair of resistance thermometers forming the other arms of each bridge. <i>C</i> , Mechanism for shifting the contact on the rheostat controlling the current in the heating coil in the final heater, shown at <i>E</i> in Plate XCIV. <i>D</i> , Temperature-difference recorder (self-balancing Wheatstone bridge) for continuously recording the difference between the temperature of the water entering and that leaving the heat absorber. <i>E</i> , "Check box" containing the ratio coils of the bridge for temperature difference measurements and coils for extending the range of differences measured, with means for checking the constancy of the resistances of the coils and the accuracy of the recorder readings and also for compensating for slight differences in the resistance of the thermometer coils when they are at the same temperature.....	720
MOTTLE-LEAF OF CITRUS TREES IN RELATION TO SOIL CONDITIONS	
PLATE H.—Various stages in mottle-leaf of the orange.....	740
PLATE XCVI.—Orange leaves showing mottle-leaf.....	740
PLATE XCVII.—A more advanced stage of mottle-leaf of orange, showing the reduction in the size of the leaves.....	740
VEGETATIVE SUCCESSION UNDER IRRIGATION	
PLATE XCVIII.—Rock Creek Valley, near Rock River Station.....	760
PLATE XCIX.—A nearer view of the bench slope; the same tree shown in Plate C, figure 1.....	760
PLATE C. Fig. 1.—Where upland and lowland meet. Fig. 2.—Characteristic draw; the stream valley beyond. Lupin, wheat-grass, white sage, and gaillardia in the foreground.....	760

	Page
PLATE CI. Fig. 1.—The bench. The course of Rock Creek is indicated by the distant trees. Fig. 2.—Part of a reservoir on the Rock Creek ranch. ....	760
PLATE CII. Fig. 1.—Lupin recessive and cat's-foot becoming dominant. Gay's sedge subphase in background. Fig. 2.—Wheat-grass phase. ....	760
PLATE CIII. Fig. 1.—Squirrel-tail phase. A few grindelias in the foreground. Fig. 2.—Rush-sedge phase (the darker areas) replacing wheat-grass phase. ....	760
PLATE CIV. Fig. 1.—Hair-grass phase. Fig. 2.—Natural meadow. ....	760
PLATE CV. Fig. 1.—Field of oats on bench. Cat's-foot and other upland plants in foreground. Fig. 2.—Alfalfa field one year after sowing. Cat's-foot and bench grasses in foreground. ....	760
AGRICULTURAL VALUE OF IMPERMEABLE SEEDS	
PLATE CVI. Fig. 1.—A row of alsike clover from impermeable seeds between two rows from permeable seeds. Fig. 2.—A row of white clover from impermeable seeds between two rows from permeable seeds. ....	796
LIFE-HISTORY STUDIES OF <i>CIRPHIS UNIPUNCTA</i> , THE TRUE ARMY WORM	
PLATE CVII.—A, Cages for rearing <i>Cirphis unipuncta</i> ; B, leaves glued together after the eggs have been deposited; C, characteristic leaves partly eaten by first-instar larvae; D, full-grown larva; E and F, characteristic pupal cells. ....	812
CONTROL OF THE POWDERY DRYROT OF WESTERN POTATOES CAUSED BY <i>FUSARIUM TRICHOETHECIOIDES</i>	
PLATE CVIII. Fig. 1.—A potato tuber infected with powdery dryrot, showing the wrinkled condition of skin due to the decay of underlying tissues. Fig. 2.—A potato tuber infected with powdery dryrot. Advanced stage. Fig. 3.—Section through a potato tuber infected with powdery dryrot, showing the internal cavities filled with the mycelium and the spores of the fungus. ....	832
THERSILOCHUS CONOTRACHELI, A PARASITE OF THE PLUM CURCULLIO	
PLATE CIX. <i>Thersilochus conotrachei</i> : Fig. 1.—Adult female. Fig. 2.—a, Adult male; b, side view of abdomen. ....	856
APHIDOLETES MERIDIONALIS, AN IMPORTANT DIPTEROUS ENEMY OF APHIDS	
PLATE CX. <i>Aphidoletes meridionalis</i> : Fig. 1.—Adult female; a, Antenna of male, showing structure; b, tip of male abdomen. Fig. 2.—Larva attacking a pea aphid. ....	888
PROGRESSIVE OXIDATION OF COLD-STORAGE BUTTER	
PLATE CXI.—Gas apparatus used in the extraction and analysis of the air confined in butter. ....	952
STUDIES ON THE PHYSIOLOGY OF REPRODUCTION IN THE DOMESTIC FOWL.— XV. DWARF EGGS	
PLATE CXII.—A collection of dwarf eggs with a normal egg in the center of the group. ....	1042
PLATE CXIII.—Fig. 1.—Ovarian follicles (b-f) and the dwarf egg a from case 27. Fig. 2.—Shell of a compound egg which was composed of two albumen masses partly separated at the level of the seam in the shell by an incomplete egg membrane. Fig. 3.—Dwarf egg containing a mass of yolk not in a yolk membrane, but separated from the albumen by a membrane-like layer of chalazal threads. Fig. 4.—Dwarf egg formed around an artificial yolk of agar which was inserted into the oviduct. a, Complete egg; b, agar yolk. ....	1042

$\alpha$ -CROTONIC ACID, A SOIL CONSTITUENT

PLATE CXIV. Fig. 1.— $\alpha$ -Crotonic acid from soil. Fig. 2.—Synthetic $\alpha$ -crotonic acid. ....	Page 1046
---	--------------

## TEXT FIGURES

CLIMATIC CONDITIONS AS RELATED TO *CERCOSPORA BETICOLA*

FIG. 1. <i>Cercospora beticola</i> : A. Section of overwintered sugar-beet leaf showing embedded sclerotia-like body, <i>a</i> , with a mass of old conidiophores, <i>b</i> , from which a new conidium, <i>c</i> , was produced. B. Production of rather typical conidiophores, <i>b</i> , and conidia, <i>c</i> , from a sclerotia-like mass, <i>a</i> , taken from overwintered host material and placed in hanging-drop cultures. ....	22
2. Curves of the maximum and minimum soil and air temperatures for the period from December 5, 1912, to March 13, 1913, at Rocky Ford, Colo., and air temperatures from December 5, 1913, to March 13, 1914, at Madison, Wis., together with the periods that snow covered the ground. ....	28
3. Curves of the maximum and minimum soil and air temperatures for Rocky Ford, Colo., from March 13 to June 17, 1913, and for Madison, Wis., from March 13 to June 17, 1914. ....	29
4. Curves of the maximum and minimum temperatures and relative humidities and the number of hours that the humidity remained above 60 from noon of the preceding to noon of the given day among sugar-beet plants and in the air 5 feet above the field, together with the field rainfall and irrigation records. June 11 to August 2, 1913, at Rocky Ford, Colo. ....	32
5. Curves of the maximum and minimum temperatures among sugar-beet plants and at the Weather Bureau station, and the seasonal rainfall records at Madison, Wis., in 1914, and the number of hours that the humidity remained above 60 among the sugar-beet plants in the field at Madison, Wis., in 1914, and at Rocky Ford, Colo., in 1913. ....	34
6. Curves of the leaf spot history series, showing the production of conidia on different dates from June 24 to September 19, 1913, at Rocky Ford, Colo. ....	48
7. Curves of the maximum and minimum temperatures and humidities, the number of hours that the humidity remained above 60 from noon of the preceding to noon of the given day among the plants, and rainfall and irrigation records, taken in a medium-early sugar-beet field from June 10 to September 22, 1913, at Rocky Ford, Colo. ....	51
8. Curves of the comparative production of conidia on the upper and lower surfaces of the leaf spots, representing series E, K, N, and G of Table V and figure 6, Rocky Ford, Colo., 1913. ....	52
9. Curves of the 2-day average increases in the number of leaf spots per plant in a medium-early and an early sugar-beet field, from June 18 to September 19, 1913, at Rocky Ford, Colo. ....	54
10. Curves of the maximum and minimum temperatures and relative humidities and the number of hours that the humidity remained above 60 from noon of the preceding to noon of the given day among the sugar-beet plants of a medium-early and an early field. August 2 to September 21, 1913, at Rocky Ford, Colo. ....	58



## CITRUS CANKER

	Page
FIG. 1. Diagrammatic representation of young open type of Citrus canker of half the diameter of the one shown in figure 2. <i>pp</i> , Palisade parenchyma; <i>ue</i> , upper epidermis; <i>le</i> , lower epidermis; <i>d</i> , diseased tissues; <i>a</i> , air space arising from tensions due to the enlargement of cells and disintegration of tissues.....	72
2. Diagrammatic representation of canker on old Citrus leaf; <i>pp</i> , Palisade parenchyma; <i>ue</i> , upper epidermis; <i>le</i> , lower epidermis; <i>p</i> , pycnidium of <i>Phoma socia</i> ; <i>d</i> , diseased tissues; <i>a</i> , air space arising from tensions due to the enlargement of cells and disintegration of tissues.....	73
<i>Pseudomonas citri</i> : <i>a</i> , Stained with carbol fuchsin; <i>b</i> , stained with Williams's flagellar stain (adapted from Hasse); <i>c</i> , stained with anilin gentian violet.....	76
4. Early stage of Citrus canker in cross section on a young leaf of seedling grapefruit.....	80
5. <i>Pseudomonas citri</i> : ( <i>a</i> ). In the mesophyll tissue and ( <i>b</i> ) in the palisade parenchyma.....	81
6. Drawing of a stained section of a natural canker on grapefruit.....	82
7. Cross section in outline of a spongy canker on the rind of a fruit of <i>Citrus decumana</i> , showing ruptured epidermis and hypertrophy of the rind tissues, the cells of which are loosely attached.....	83
8. <i>a</i> , Cross section of a pycnidium of <i>Phoma socia</i> from a grapefruit leaf. <i>b</i> , Germination of conidia of <i>Phoma socia</i> after 24 hours in water. <i>c</i> , Mycelium of this fungus in old cultures.....	86

## DETERMINATION OF STEARIC ACID IN BUTTER FAT

FIG. 1. Exterior of constant-temperature crystallization tank.....	104
2. Interior of constant-temperature crystallization tank.....	105

## LIFE HISTORY AND HABITS OF TWO NEW NEMATODES PARASITIC ON INSECTS

FIG. 1. <i>Diplogaster labialis</i> : <i>A</i> , Mating; <i>B</i> , mature female reared in water culture, <i>a</i> , lip region, <i>b</i> , esophagus, <i>c</i> , median bulb, <i>d</i> , cardiac bulb, <i>e</i> , intestine, <i>f</i> , ovaries. <i>g</i> , egg, <i>h</i> , genital pore, <i>i</i> , rectum, <i>k</i> , anus; <i>C</i> , mature male reared in water culture, <i>a</i> , lip region, <i>b</i> , esophagus, <i>c</i> , median bulb, <i>d</i> , cardiac bulb, <i>e</i> , intestine, <i>k</i> , anus, <i>m</i> , spicula; <i>D</i> , at time of hatching; <i>E</i> , female during process of molting; <i>F</i> , dead female with young nematode which hatched within her body.....	116
2. <i>A-H</i> , <i>Diplogaster labialis</i> : Development of the egg; <i>I</i> , <i>Diplogaster aerivora</i> : mature male reared in moist soil; <i>J</i> , <i>Diplogaster aerivora</i> : mature male reared in water culture, <i>a</i> , lip region, <i>b</i> , esophagus, <i>c</i> , median bulb, <i>d</i> , cardiac bulb, <i>e</i> , intestine, <i>k</i> , anus, <i>m</i> , spicula; <i>K</i> , <i>Diplogaster aerivora</i> : dead female with young which hatched within her body; <i>L</i> , <i>Diplogaster aerivora</i> : mating.....	117
3. <i>Diplogaster aerivora</i> : <i>A</i> , Form found in termite; <i>B</i> , at time of hatching; <i>C</i> , female reared in water culture, not quite mature, <i>a</i> , lip region, <i>b</i> , esophagus, <i>c</i> , median bulb, <i>d</i> , cardiac bulb, <i>e</i> , intestine, <i>f</i> , ovaries, <i>h</i> , genital pore, <i>i</i> , rectum, <i>k</i> , anus; <i>D</i> , mature female reared in moist soil; <i>E</i> , mature female reared in water culture; <i>F-M</i> , development of the egg.....	122

## EFFECT OF PASTEURIZATION ON MOLD SPORES

FIG. 1. Curve of the number of species of molds surviving pasteurization of milk for 30 minutes at a series of temperatures.....	157
2. Curve of the number of species of molds surviving flash pasteurization at a series of temperatures.....	160

	Page
FIG. 3. Curve of the number of species of molds surviving dry heat for 30 minutes at a series of temperatures. ....	164
EFFECT OF WATER IN THE RATION ON THE COMPOSITION OF MILK	
FIG. 1. Grouping of cow and kind of ration fed cows 23, 24, 25, and 27. ....	173
HYPERASPIS BINOTATA, A PREDATORY ENEMY OF THE TERRAPIN SCALE	
FIG. 1. Map showing the distribution in the United States of <i>Hyperaspis binotata</i> . ....	198
TESTS OF THREE LARGE-SIZED REINFORCED-CONCRETE SLABS UNDER CONCENTRATED LOADING	
FIG. 1. Diagram illustrating the method of obtaining "effective width" in reinforced-concrete slab tests. ....	207
2. Diagram showing location of strain-gauge points on top of slab 835. ....	211
3. Diagram showing location of strain-gauge points on top of slab 930. ....	212
4. Diagram showing location of strain-gauge points on top and bottom of slab 934. ....	213
5. Concrete deformation curves for concentrated center load on slab 835. ....	214
6. Concrete deformation curves for slab 930. ....	215
7. Deformation curves for slab 934. ....	216
8. Deformation curves for slab 934, computed from first zero reading. ....	216
9. Concrete deformation curves for slab 835 with 2-point loading. ....	217
10. Concrete deformation curves for slab 934 with 2-point loading. ....	217
11. Steel deformation curves for slab 934 with 2-point loading. ....	218
12. Concrete deformation curves for slab 835 with 4-point loading. ....	219
13. Concrete deformation curves for slab 934 with 4-point loading. ....	219
14. Deflection curves for slab 934 on first application of load. ....	220
15. Deflection curves for slab 934 on second application of load. ....	221
16. Deflection curves for slab 934 with 2-point loading. ....	222
17. Diagram showing effect of breaking load on slab 835. ....	223
18. Diagram showing effect of breaking load on slab 930. ....	223
19. Diagram showing effect of breaking load on slab 934. ....	224
20. Concrete deformation curves for a 10,000-pound concentrated center load on slab 934. Variation in deformations plotted parallel to supports. ....	225
21. Concrete deformation curves for a 10,000-pound concentrated center load on slab 934. Variation in deformations plotted perpendicular to supports. ....	226
22. Concrete deformation curves for slab 934. Lateral deformations plotted both parallel and perpendicular to supports. ....	227
23. Iso-deformation lines for slab 934 under concentrated center load. ....	228
24. Concrete deformation curves for slab 934 under 40,000-pound 4-point loading. Deformations measured perpendicular to supports. Variation of deformations plotted parallel to the supports. ....	229
25. Concrete deformation curves for slab 934 under 40,000-pound 4-point loading. Deformations measured perpendicular to supports. Variation of deformations plotted perpendicular to supports. ....	230
26. Iso-deformation lines for slab 934 under 40,000-pound 4-point loading. Deformations measured perpendicular to supports. ....	231
27. Curves showing effective width versus load (concentrated center load). ....	231
28. Curve showing effective width versus thickness. ....	232

## EFFECT OF CLIMATIC FACTORS ON THE HYDROCYANIC-ACID CONTENT OF SORGHUM

	Page
FIG. 1. Curves showing the hydrocyanic-acid content of sorghum on the various plots.....	264
2. Curves showing the rate of growth of the sorghum on the various plots..	265
3. Curves showing the precipitation, temperature, and humidity relations at the various experiment stations during the growing season of 1915..	266
4. Curves showing the contemporary climatic conditions at the various plots, together with crop data and hydrocyanic-acid content.....	267

## HYPODERMA DEFORMANS, AN UNDESCRIBED NEEDLE FUNGUS OF THE WESTERN YELLOW PINE

FIG. 1. A side view of two apothecia of <i>Hypoderma deformans</i> on needles of <i>Pinus ponderosa</i> , showing the longitudinal medial split.....	278
2. Asci, spores, and paraphyses of <i>Hypoderma deformans</i> .....	279
3. Cross section of an apothecium of <i>Hypoderma deformans</i> on a needle of <i>Pinus ponderosa</i> , showing mature asci with spores, the point of first rupture, and the tissues of the leaf most seriously affected by the mycelium of the fungus.....	280
4. The upper portion of a young ascus of <i>Hypoderma deformans</i> , showing the formation of the pore at the tip through which the spores are expelled.....	281

## COMPARATIVE STUDY OF THE ROOT SYSTEMS AND LEAF AREAS OF CORN AND THE SORGHUMS

FIG. 1. Evaporation from a free water surface (tank) at Garden City, Kans., during the growing seasons of 1914 and 1915.....	314
2. Comparison of the leaf areas of Pride of Saline corn, Black-hull kafir, and Dwarf milo at four stages of the growth of these plants during the season of 1914.....	328
3. A graphic illustration of the sheath areas of Pride of Saline corn, Black-hull kafir, and Dwarf milo at four stages of the growth of these plants during the season of 1914.....	329

## WOOLLY PEAR APHIS

FIG. 1. Comparative structure of antennae and wax pores of <i>Eriosoma</i> spp.: A, distal segments of antenna of winged viviparous female of <i>E. pyricola</i> ; B, distal segments of antenna of winged viviparous female of <i>E. ulmi</i> ; C, distal segments of antenna of wingless viviparous female of <i>E. americanum</i> ; D, distal segments of antenna of wingless viviparous female of <i>E. lanigerum</i> ; E, distal segments of antenna of wingless viviparous female of <i>E. pyricola</i> ; F, distal segments of antenna of winged viviparous female of <i>E. americanum</i> ; G, distal segments of antenna of winged viviparous female of <i>E. lanigerum</i> ; H, compound wax pore of <i>E. lanigerum</i> ; I, compound wax pore of <i>E. pyricola</i> ; J, distal segments of antenna of first instar wingless viviparous female of <i>E. pyricola</i> .....	359
--	-----

## STIMULATING INFLUENCE OF ARSENIC UPON THE NITROGEN-FIXING ORGANISMS OF THE SOIL

FIG. 1. Graph showing the action of five compounds of arsenic on nitrogen fixation in dry soil.....	302
2. Graph showing the effect of aeration on the nitrogen-fixing activity of soil containing compounds of arsenic.....	408

	Page
FIG. 3. Graph showing the effect of heat on the nitrogen-fixing power of soil treated and not treated with arsenic.....	409
4. Effect of various arsenic compounds in the ratio of 400 parts of the compound to 1,000,000 parts of soil on the activity of various soil organisms.....	411
5. Graph showing parts per million of various arsenic compounds in the soil at which the greatest stimulation occurred.....	412
TRANSMISSION AND CONTROL OF BACTERIAL WILT OF CUCURBITS	
FIG. 1. Comparison of the amount of wilt with striped-beetle prevalence and with meteorological phenomena in three fields, East Marion, Long Island, N. Y., season of 1915.....	421
2. Comparison of relative wilt control of Bordeaux mixture plus lead arsenate, Bordeaux mixture alone, and lead arsenate alone in field 1, East Marion, Long Island, season of 1915.....	430
3. Curves showing relative wilt control of Bordeaux mixture and lead arsenate with date of first application as a variant in field 1, East Marion, Long Island, season of 1915.....	431
ALEYROBIDAE, OR WHITE FLIES, ATTACKING THE ORANGE, WITH DESCRIPTION OF THREE NEW SPECIES OF ECONOMIC IMPORTANCE	
FIG. 1. <i>Aleurocanthus citripertus</i> : A, Pupa case; B, egg; C, polygonal markings of egg; D, vasiform orifice of pupa case; E, spine from dorsum of pupa case; F, margin of pupa case; G, genitalia of adult male; H, forewing of male; I, antenna of pupa case; J, leg of pupa case; K, L, marginal teeth, much enlarged; M, central swollen spine from dorsal area.....	460
2. <i>Aleurocanthus woglumi</i> : A, Egg; B, polygonal markings of egg; C, pupa case; D, margin of pupa case; E, vasiform orifice of pupa case; F, forewing of adult female; G, same, showing variation in markings; H, costal margin at base of wing of female; I, forewing of male; J, male genitalia.....	462
3. <i>Aleurothrixus porteri</i> , <i>A. howardi</i> , and <i>A. floccosus</i> : A, <i>Aleurothrixus porteri</i> : Larva, first instar. B, <i>A. porteri</i> : Caudal spine of pupa case. C, <i>A. porteri</i> : Clasper of male. D, <i>A. porteri</i> : Egg. E, <i>A. howardi</i> : Caudal spine. F, <i>A. porteri</i> : Pupa case. G, <i>A. porteri</i> : Forewing of adult. H, <i>A. floccosus</i> : Vasiform orifice of pupa case. I, <i>A. porteri</i> : Vasiform orifice of pupa case. J, <i>A. howardi</i> : Vasiform orifice of pupa case. K, <i>A. porteri</i> : Margin of pupa case. L, <i>A. porteri</i> : Margin of early larva.....	467
RELATIVE WATER REQUIREMENT OF CORN AND THE SORGHUMS	
FIG. 1. Curves of the evaporation at Garden City, Kans., for the growing period of 1915.....	474
CALIFORNIA GREEN LACEWING FLY	
FIG. 1. The California green lacewing fly: Adult.....	515
2. <i>Chrysopa californica</i> : Eggs.....	518
3. <i>Chrysopa californica</i> : First instar.....	519
4. <i>Chrysopa californica</i> : Third instar.....	521
5. <i>Chrysopa californica</i> : Pupal case.....	522
6. <i>Chrysopa californica</i> : Pupa.....	522
7. <i>Chrysopa californica</i> : Pupa freshly emerged from its cocoon.....	523

## STORAGE-ROTS OF ECONOMIC AROIDS

- FIG. 1. Spores of different storage-rot organisms: A, *Diplodia tubericola* from dasheen; B, *Diplodia tubericola* from sweet potato; C, *Diplodia gossypina* from cotton; D, *Diplodia* sp. from *Mungifera indica*; E, *Diplodia maculata* from *Toxylon pomiferum*; F, *Fusarium solani*. . . . . 552

## LARVAL CHARACTERS AND DISTRIBUTION OF TWO SPECIES OF DIATRAEA

- FIG. 1. a, Average angle formed by imaginary lines through bases of setae of *Diatraea saccharalis crambidoides*; b, average angle formed by imaginary lines through bases of setae of *D. zeacollla*. . . . . 622

## THE DISEASE OF POTATOES KNOWN AS "LEAK"

- FIG. 1. Microscopical appearance of *Pythium debaryanum* isolated from potatoes affected with potato leak: a, Cell of a potato tuber showing fungus filaments therein; b, oogonia and antheridia; c, mycelium; d, germinating conidia. . . . . 630

## LIFE CYCLES OF THE BACTERIA

- FIG. 1. Life cycle of *Bacillus azotobacter*. The broken straight lines divide the different types of growth indicated by the letters A to M. The Greek letters  $\alpha$  to  $\lambda$  refer to subdivisions. The single and double pointed arrows indicate the development of one form from another. The four circles confine, in every case, all those forms which represent together a rather constant mode of life and which have been usually considered as bases for establishing separate species. . . . . 678

## MOTTLE-LEAF OF CITRUS TREES IN RELATION TO SOIL CONDITIONS

- FIG. 1. Graphical presentation of the relationship between humus content of soil and percentage of mottled orange leaves. . . . . 731
2. Graphical presentation of the relationship between the ratio of organic carbon to humus in the soil and the percentage of mottled orange leaves. . . . . 734
3. Graphical presentation of the relationship between the ratio of humus to lime in the soil and the percentage of mottled leaves. . . . . 735
4. Graphical presentation of the relationship between the mineral carbonates in the soil and the percentage of mottled lemon leaves. . . . . 737

## AGRICULTURAL VALUE OF IMPERMEABLE SEEDS

- FIG. 1. Curves showing the rate of softening of impermeable red-clover seeds of different degrees of maturity. . . . . 767
2. Curves showing the rate of softening and of germination of impermeable red-clover seeds of different lots. . . . . 768
3. Curves showing the changes in the permeability of seeds in wet blotters and in dry storage for various periods. . . . . 773
4. Curves showing the rate of softening of impermeable red-clover seeds under different temperature conditions. . . . . 780
5. Curves of the seedling production in the field in 16 to 18 days and of the germination in chamber in 8 days. . . . . 787
6. Curves of the seedling production in the field and of the germination in chamber in 12½ months. . . . . 790

## LIFE-HISTORY STUDIES OF CIRPHIS UNIPUNCTA, THE TRUE ARMY WORM

	Page
FIG. 1. Diagram of relative amounts of foliage eaten in each larval instar by <i>Cirphis unipuncta</i> .....	861
2. Posterior extremity of male and female pupa: <i>a</i> , Male; <i>b</i> , female.....	812

## THERSILOCHUS CONOTRACHELI, A PARASITE OF THE PLUM CURCULIO

FIG. 1. <i>Thersilochus conotrachei</i> : Cocoon.....	852
2. <i>Thersilochus conotrachei</i> : Egg.....	852
3. <i>Thersilochus conotrachei</i> : Larva of first instar.....	852
4. <i>Thersilochus conotrachei</i> : Ventral surface of head of first larval instar: <i>a</i> , Labrum; <i>b</i> , maxilla; <i>c</i> , labium; <i>d</i> , mandible.....	853
5. <i>Thersilochus conotrachei</i> : Full-grown larva of first instar.....	853
6. <i>Thersilochus conotrachei</i> : Larva of second instar.....	853
7. <i>Thersilochus conotrachei</i> : Full-grown larva of fifth instar.....	854
8. <i>Thersilochus conotrachei</i> : Face of full-grown larva: <i>a</i> , Labrum; <i>b</i> , mandible; <i>c</i> , maxilla; <i>d</i> , maxillary palpus; <i>e</i> , labium; <i>f</i> , labial palpus ..	854
9. <i>Thersilochus conotrachei</i> : Pupa of female, and apex of abdomen of male pupa.....	854

## EFFECT ON PLANT GROWTH OF SODIUM SALTS IN THE SOIL.

FIG. 1. Diagram of the percentage of sodium carbonate added to the soil in experiment 1 (1913), with the percentage of carbonate and bicarbonate recovered and the total green weight of wheat obtained.....	860
2. Diagram of the percentage of sodium carbonate added to the soil in experiment 2 (1914), with the percentage of carbonate and bicarbonate recovered and the green weight of wheat obtained.....	862
3. Diagram of the percentage of sodium carbonate added to the loam soil in experiment 3 (1915), with the percentage of carbonate and bicarbonate recovered and green weight of wheat.....	864
4. Diagram of the percentage of sodium carbonate added to Monterey sand in experiment 3 (1915), with the percentage of carbonate and bicarbonate recovered and the total green weight of wheat obtained.....	864
5. Diagram of the percentage of sodium bicarbonate added to the soil in experiment 4 (1914), with carbonate and bicarbonate recovered, together with the total green weight of wheat obtained.....	865
6. Diagram of the decrease in growth of wheat seedlings in experiments 2 and 4 as affected by the total carbonate salts recoverable from the soil.....	866
7. Diagram of the quantity of sodium chlorid added to the soil, with the quantity of chlorid recovered, and the total green weight of wheat obtained.....	867
8. Diagram of the quantity of sodium sulphate added to the soil in experiment 6, the quantity recovered, and the total green weight of wheat obtained.....	868

## APHIDOLETES MERIDIONALIS, AN IMPORTANT DIPTEROUS ENEMY OF APHIDS

FIG. 1. <i>Aphidoletes meridionalis</i> : Eggs in situ on leaf of rape: <i>a</i> , egg.....	884
2. <i>Aphidoletes meridionalis</i> : Larva, dorsal view.....	886
3. <i>Aphidoletes meridionalis</i> : <i>a</i> , Cocoon formed on surface of ground; <i>b</i> , cocoon formed on a leaf blade.....	886
4. <i>Aphidoletes meridionalis</i> : Pupa, lateral view.....	887

## INFLUENCE OF BARNYARD MANURE AND WATER UPON THE BACTERIAL ACTIVITIES OF THE SOIL.

	Page
Fig. 1. Curves of the ammonifying powers of soil in pots with varying quantities of manure and water.....	899
2. Curves of the nitrifying powers of soil in pots with varying quantities of manure and water.....	900
3. Curves of the number of colonies of bacteria developing from fallow soil with varying quantities of manure and water.....	904
4. Curves of the ammonifying powers of fallow soil with varying quantities of manure and water.....	907
5. Curves of the nitrifying powers of fallow soil with varying quantities of manure and water.....	910
6. Curves of the number of colonies of bacteria developing from cropped plats with varying quantities of manure.....	912
7. Curves of the ammonifying powers of soil of cropped plats with varying quantities of manure and water.....	915
8. Curves of the nitrifying powers of soil of cropped plats with varying quantities of manure and water.....	918
9. Diagram of the influence of manure on the yield and bacterial activities of a soil.....	920
10. Diagram of the influence of irrigation water on the yield and bacterial activities of a soil.....	921

## PROGRESSIVE OXIDATION OF COLD-STORAGE BUTTER

Fig. 1. Diagram of gas apparatus used in the extraction and analysis of the air confined in butter.....	931
---	-----

## BACTERIOLOGICAL STUDIES OF A SOIL, SUBJECTED TO DIFFERENT SYSTEMS OF CROPPING FOR 25 YEARS

Fig. 1. Curves of the nitrate formation in soil of fertility plats at Columbia, Mo., in 1913-14, after 28 days' incubation with the addition of 60 mgm. of nitrogen as cottonseed meal, but without the addition of calcium carbonate.....	962
--	-----

STUDIES ON THE PHYSIOLOGY OF REPRODUCTION IN THE DOMESTIC FOWL.—  
XV. DWARF EGGS

Fig. 1. Diagram showing the percentage of the yearly total egg production (8-year average, 1899-1907) and the percentage of the total dwarf-egg production (8-year average, 1908-1916) which occurred during each month.....	1004
2. Diagram showing for the years 1911-12 and 1914-15 combined the percentage of the yearly total egg production and dwarf-egg production which occurred during each month and 100 times the percentage of the eggs produced each month which were dwarf.....	1005
3. Diagram showing the number of dwarf eggs which occurred in each tenth of a litter.....	1011
4. Diagram showing the number of dwarf eggs which occurred in each fifth of a litter.....	1014

# JOURNAL OF AGRICULTURAL RESEARCH

DEPARTMENT OF AGRICULTURE

VOL. VI

WASHINGTON, D. C., APRIL 3, 1916

NO. 1

## RELATION OF CARBON BISULPHID TO SOIL ORGANISMS AND PLANT GROWTH<sup>1</sup>

By E. B. FRED,

*Agricultural Bacteriologist, Agricultural Experiment Station  
of the University of Wisconsin*

### INTRODUCTION

In a previous publication concerning the action of carbon bisulphid ( $\text{CS}_2$ ) on bacteria and plants data were presented to show the beneficial effect of this substance on the soil flora (1).<sup>2</sup> The increased plant growth following the addition of carbon bisulphid in many cases is enormous. For example, a small application often causes an increase in yield from 100 to 200 per cent. It is impossible to account for this remarkable gain on the assumption that the only action of the carbon bisulphid is that of added plant food. It was found, as has been noted by many investigators (5, 6, 11, 12), that this volatile antiseptic exerts a very decided effect on the micro-organisms of the soil. As measured by plate counts, there is at first usually a great decrease in numbers, followed by a period of excessive increase, the total numbers far exceeding those that ordinarily exist. In certain cases carbon bisulphid has not only failed to cause an increase in plant growth, but has, on the contrary, caused a decrease.

Search has been made by many investigators for a satisfactory explanation of this peculiar action of carbon bisulphid. Many theories have been advanced. Concerning these theories so much has been written that a detailed discussion of the literature seems unnecessary. Indeed, it would be impossible within the limited scope of this paper to present a summary of the various explanations. One point is very prominent in nearly all of the publications: The action of carbon bisulphid is varied. Because of the interest attached to this problem, it was arranged to study some of the factors that might influence the action of carbon bisulphid. The experiments described in this paper are discussed under three main heads: First, the effect of varying amounts of carbon bisulphid; second, the effect of carbon bisulphid on various plants; and third, the effect of carbon bisulphid in various soils. In all of this work fresh field soil and commercial carbon bisulphid were used. Some of the experiments represent a combined study of the effect on both the lower and higher forms of plant life.

<sup>1</sup> Published with permission of the Director of the Wisconsin Agricultural Experiment Station.

<sup>2</sup> Reference is made by number to "Literature cited," p. 12-16.



## EXPERIMENTAL METHODS

Commercial carbon bisulphid was poured into small holes in the soil, and these were covered immediately. The soil was sieved and potted in 2-gallon jars and the moisture maintained at half saturation. Changes in the soil flora were determined at regular intervals by plate counts of the number of bacteria and dilution counts of the number of active protozoa. The formation of ammonia and nitrates was measured at regular intervals.

The following plants were used: Buckwheat (*Fagopyrum fagopyrum*), clover (*Trifolium pratense*), corn (*Zea mays*), mustard (*Sinapis alba*), oats (*Avena sativa*), and rape (*Brassica napus*). In many of the experiments a first and a second crop were grown.

## EFFECT OF CARBON BISULPHID ON THE NUMBER AND ACTIVITY OF SOIL ORGANISMS

Eight jars were filled with Miami silt-loam soil from the Experiment Station farm. These were arranged in duplicate and treated as follows: (1) Control, untreated; (2) 2 per cent of carbon bisulphid; (3) 2 per cent of carbon bisulphid, evaporated; (4) 2 per cent of carbon bisulphid, evaporated, and reinoculated with 5 per cent of the original soil.

Twenty-four hours after treatment the soil in the evaporated series was spread out on sterile paper and the volatile antiseptic allowed to escape. At the end of the second 24-hour period the soil was put back into the jars. In order to prevent any contamination, the jars were covered with a double layer of cheesecloth and nonabsorbent cotton. This cover should allow free access of air without much danger of contamination. At regular intervals the covers were removed and samples drawn for analysis. The results of these determinations are presented in Tables I and II.

## NUMBER OF ORGANISMS

BACTERIA.—In Table I are shown the number of bacteria in 1 gm. of soil at different times and under the different conditions.

TABLE I.—Effect of carbon bisulphid on number of bacteria

Time.	Bacteria per gram of dry soil.			
	Control.	2 per cent of carbon bisulphid.	2 per cent of carbon bisulphid evaporated.	2 per cent of carbon bisulphid evaporated + 5 per cent of soil from control.
<i>Days.</i>				
1.....	17,496,000	1,965,000	2,260,000	2,358,000
3.....	22,010,000	23,975,000	8,254,000	12,489,000
5.....	20,635,000	25,253,000	27,416,000	95,499,000
9.....	14,739,000	36,651,000	61,904,000	.....
13.....	16,115,000	90,473,000	98,850,000	80,420,000
21.....	19,508,000	60,149,000	71,257,000	52,495,000
25.....	18,272,000	68,276,000	86,483,000	64,570,000
29.....	15,340,000	90,645,000	84,272,000	38,495,000
60.....	12,372,000	58,101,000	60,000,000	30,000,000

At first the antiseptic causes a great reduction in the number of organisms capable of developing on Heyden agar. The period of depression lasts for only a short time—in this experiment about five days. From that time until the end of the test the number of organisms in the treated series far exceeded that of the control. The highest number in the carbon bisulphid evaporated and unevaporated soil occurred about the thirteenth day; while the carbon bisulphid evaporated soil plus control soil gave the highest count on the fifth day. At the time of the last count, 60 days after carbon bisulphid was added, the organisms in the treated series far exceeded those in the original soil. Apparently the effect of carbon bisulphid on the number of bacteria is noticeable for a long period of time.

If the results of the counts with carbon bisulphid unevaporated are compared with those of carbon bisulphid evaporated, it appears that no very marked difference exists. The greatest reduction in numbers occurred in soils with the carbon bisulphid evaporated. It is significant that soil with carbon bisulphid evaporated should prove more injurious to micro-organisms than the unevaporated. This agrees with Gainey (2, p. 592), who reports that the combined effect of the two processes seemed more injurious to nitrification than treatment with carbon bisulphid unevaporated.

After the thirteenth day the treated and reinoculated soil did not show as many organisms as the treated series. This difference is shown very distinctly in Plate I, which is reproduced from a photograph of a number of colonies developing on agar. Four parallel plates were made from the same dilution of each soil.

On this date samples were also drawn for ammonification tests. The purpose of this was to measure the rate of the decomposition of casein in the various series, and 1 per cent of casein was added to the soil and the ammonia determined after 12 and 24 hours. The beneficial effect of carbon bisulphid on ammonification is very evident. If after 12 hours the untreated is 100, then carbon bisulphid unevaporated is 154, carbon bisulphid evaporated is 212, and carbon bisulphid reinoculated is 190.

After 24 hours the untreated is equal to 100, carbon bisulphid unevaporated is 149, carbon bisulphid evaporated is 171, and carbon bisulphid reinoculated is 153. The data show very clearly that casein is decomposed more rapidly in treated than in untreated soils. This difference is most prominent in the 12-hour tests.

PROTOZOA.—Counts at the beginning showed the presence of protozoa in dilutions representing 1 to 1,000 gm. of soil (13, p. 626). Two weeks after treatment the soils were recounted. At this time numerous small flagellates were found in dilutions of 1 to 1,000. It is evident that the different treatments with carbon bisulphid had not seriously injured this group of organisms.

**AZOTOBACTER.**—One month after treatment with carbon bisulphid, qualitative tests were made. The Azotobacter organisms were found in all soils. The brown film of Azotobacter from the treated soils was not so profuse as that from the original soil.

**ALGÆ.**—In order to estimate the number of algæ, dilution tests were made. These cultures were incubated for 30 days. The smaller forms were found in great numbers in all of the soils.

The important facts in these data are (1) that the volatile antiseptic fails to remove these larger soil organisms and (2) that the smaller forms of bacteria are only temporarily reduced. The decrease in numbers is soon followed by a period of excessive growth.

#### ACTIVITY OF ORGANISMS

A rapid multiplication of bacteria should naturally be followed by a parallel increase in decomposition products. Accordingly samples for analysis were drawn from the jars used in the previous experiment. The results of these periodic analyses are presented in Table II.

TABLE II.—*Effect of carbon bisulphid on ammonia and nitrate content of soil*

Time.	Nitrogen per 100 gm. of dry soil.							
	Ammonia.				Nitrate.			
	Control.	2 per cent of carbon bisulphid.	2 per cent of carbon bisulphid evaporated.	2 per cent of carbon bisulphid evaporated + 5 per cent of soil from control.	Control.	2 per cent of carbon bisulphid.	2 per cent of carbon bisulphid evaporated.	2 per cent of carbon bisulphid evaporated + 5 per cent of soil from control.
Days.	Mgm.	Mgm.	Mgm.	Mgm.	Mgm.	Mgm.	Mgm.	Mgm.
At beginning	1.60	1.60	1.60	1.60	2.66	2.66	2.66	2.66
30. ....	1.68	5.27	5.41	4.71	3.35	2.50	2.00	5.55
45. ....	2.38	8.40	7.70	4.90	3.75	2.70	2.50	5.66
60. ....	2.59	5.43	5.32	2.31	4.00	2.81	2.50	4.50
75. ....	1.85	5.60	.....	2.10	3.20	2.40	2.60	5.00
90. ....	2.94	4.06	4.06	2.24	4.00	5.00	3.32	6.66

In the soils treated with carbon bisulphid there is a very decided accumulation of ammonia nitrogen. If the figures of Table I are compared with those of Table II, ammonia production, it will be seen that an increase in the number of bacteria within a certain range results in a gain in ammonia. After 30 days the amount of ammonia nitrogen in the treated soils averaged more than three times that in the original soil. After 60 days the ammonia content in the carbon bisulphid and carbon bisulphid evaporated soil was about double that of the control, while in the carbon bisulphid evaporated plus 5 per cent fresh soil it was

less. From the data it appears that reinoculation prevents large accumulations of ammonia. This is no doubt due to the oxidation of ammonia by the nitrifying bacteria. The figures of the last column (nitrate accumulation) support this statement. A stimulation of ammonification is still noticeable at the end of 3 months.

The nitrate-forming bacteria apparently do not recover so rapidly from carbon bisulphid treatment as the ammonia-producing organisms; consequently, there is no increase in nitrates until the end of 3 months. An exception to this is noted in the reinoculated soil. Here the activity of the nitrifying bacteria is evident 30 days after inoculation.

In order to ascertain, as nearly as possible, the effect of carbon bisulphid on the soluble nitrogen of the soil, the figures of Table II, ammonia and nitrate nitrogen, were combined in Table III.

TABLE III.—*Effect of carbon bisulphid on soluble nitrogen*

Time.	Ammonia and nitrate nitrogen per 100 gm. of dry soil.			
	Control	2 per cent carbon bisulphid.	2 per cent carbon bisulphid evaporated.	2 per cent carbon bisulphid evaporated + 5 per cent of soil from control.
Days.	Mgm.	Mgm.	Mgm.	Mgm.
At beginning.....	4.26	4.26	4.26	4.26
30.....	5.03	8.47	7.41	10.26
45.....	5.13	11.10	10.20	10.56
60.....	6.50	8.24	7.82	6.87
75.....	5.05	8.00	.....	7.10
90.....	6.94	9.06	7.38	8.90

From the data in this table it is very evident that carbon bisulphid causes a large increase in ammonia and nitrate nitrogen. There seems to be very little difference between the effect of the various treatments of carbon bisulphid on the formation of ammonia and nitrate nitrogen. When compared with the control soil, it will be seen that 45 days after treatment the carbon-bisulphid soils contain more than twice as much soluble nitrogen. The higher ammonia and nitrate content is very marked 90 days after treatment. A repetition of this experiment gave similar results.

A review of the data in Tables II and III shows very clearly that carbon bisulphid in Miami soil increases the total soluble nitrogen—namely, ammonia and nitrates. One interesting fact that appears from a comparison of the ammonia and nitrate content is that these two substances are to a certain degree inversely proportional.

## EFFECT OF CARBON BISULPHID ON THE HIGHER AND LOWER FORMS OF PLANT LIFE

From the results of the preceding experiments it seems that carbon bisulphid should exert a beneficial effect on the growth of higher plants. At first this should be most marked with ammonia-feeding plants, and later with nitrate-feeding plants. Unfortunately it is not possible to secure plants that feed entirely on nitrates or ammonia. For this reason it was thought best to study the relation of carbon bisulphid to the growth of several different plants. Accordingly a combination study of the effect of carbon bisulphid on higher plants and on bacteria was made. A wide range of soil types, as well as different higher plants, was used.

Before entering upon a study of the relation of carbon bisulphid to soil type and various plants, it was desired to obtain some idea of the influence of various amounts of carbon bisulphid on plant growth. The procedure was as follows: Ten kgm. of field soil (Miami silt loam) were placed in each of sixteen 2-gallon jars. The carbon bisulphid was added in varying amounts, from 0.5 per cent to 2 per cent. It was poured into holes in the soil. These holes were closed immediately and the water increased to half saturation. In order to overcome the injurious effect of carbon bisulphid, the jars were then allowed to stand for two weeks before planting.

## CORN AND MUSTARD IN MIAMI SILT LOAM

The results of the test with corn and mustard are given in Table IV. It is evident from the data of the table that these plants do not respond alike to carbon bisulphid.

TABLE IV.—Effect of varying amounts of carbon bisulphid on the growth of corn and mustard

No.	Soil	Carbon bisulphid added.	Weight of corn.			Weight of mustard.		
			Green.	Dry.	Average.	Green.	Dry.	Average.
		Per cent.	Gm.	Gm.	Gm.	Gm.	Gm.	Gm.
1.	Miami.	Control.	75	20	22.5	58	9.5	9.25
2.	do.	Control.	80	25		49	9	
3.	do.	0.5	82	18	19.5	69	13	12.50
4.	do.	.5	83	21		72	12	
5.	do.	1	132	28	19.5	95	13	12.50
6.	do.	1	22	11		75	12	
7.	do.	2	85	21	24	105	16	16.50
8.	do.	2	125	27		112	17	

In all concentrations except 2 per cent, carbon bisulphid injured the growth of corn. Mustard, on the other hand, was greatly benefited by the carbon-bisulphid treatment. An increased growth was observed from all concentrations. The maximum gain was noted with 2 per cent of carbon bisulphid. This beneficial effect on mustard is very evident from Plate II, figure 1. If this increase in growth is due to the larger

amount of soluble nitrogen as ammonia or nitrate, then corn and mustard should behave much alike. The nitrogen-feeding power of these plants has been studied by Krüger (8), Gerlach and Vogel (3), and others. It is supposed that both corn and mustard are heavy nitrogen-feeding crops, able to take nitrogen either in the form of ammonia or nitrate.

#### BUCKWHEAT, CORN, AND OATS IN MIAMI SILT LOAM

In order to decrease the factor of individual variation, four parallel jars of Miami silt loam were used in each series in the following experiment. For the second crop these were subdivided into sets of two each. After the first crop was harvested, the soil and roots were thoroughly mixed and the jars replanted. The rotation was as follows: First crop, buckwheat; second crops, corn and mustard; first crop, corn; second crop, buckwheat; first crop, oats; second crops, corn and mustard. In Tables V, VI, and VII are presented the results of these experiments.

TABLE V.—*Effect of carbon bisulphid on the growth of buckwheat and corn*

No.	Soil.	Carbon bisulphid added.	Weight of first crop, buckwheat.			Weight of second crop, corn.		
			Green.	Dry.	Average.	Green.	Dry.	Average.
1	Miami	Per cent. Control.	Gm.	Gm.	Gm.	Gm.	Gm.	Gm.
2	do.	Control.	90	15.5	19	152	28.5	31
3	do.	Control.	97	18		160	33.5	
4	do.	Control.	121	22.2				
5	do.	Control.	126	20.5				
6	do.	2	124	23	24.5	160	34	32.7
7	do.	2	145	26.5		150	31.5	
8	do.	2	127	23				
	do.	2	126	25.5				

The yields of buckwheat and corn are given in Table V. The weights of the mustard were lost. Buckwheat gave an increase in the treated soil, while corn (the second crop) did not show any improvement. Determinations of ammonia present at the time the buckwheat was cut (three months after treatment) resulted as follows: Ammonia—if control is 100, then carbon bisulphid treated is 192. Nitrate—if control is 100, then carbon bisulphid treated is 28. The antiseptic increases ammonia, but decreases the nitrate content of soil. The results of investigation show that buckwheat feeds largely on nitrate nitrogen (9), while corn is supposed to be able to take its nitrogen in the form of ammonia. A difference in nitrogen-feeding power can not be used to explain the unequal behavior of these plants toward carbon bisulphid. Although the weights of the mustard crop were not kept, the action of the carbon bisulphid was evident. There was a decided gain in the growth of plants in the treated series.

From the data of Table VI it is obvious that carbon bisulphid has very little effect on corn (first crop) or buckwheat (second crop).

TABLE VI.—Effect of carbon bisulphid on the growth of corn and buckwheat

No.	Soil.	Carbon bisulphid added.	Weight of first crop, corn.			Weight of second crop, buckwheat.		
			Green.	Dry.	Average.	Green.	Dry.	Average.
1.	Miami.	Per cent.	Gm.	Gm.	Gm.	Gm.	Gm.	Gm.
2.	do.	Control.	480	85	84.7	115	26	23.5
3.	do.	Control.	440	82		125	27	
4.	do.	Control.	500	90		100	22	
5.	do.	Control.	410	82		95	19	
6.	do.	2	380	77	83	128	27	20.5
7.	do.	2	400	85		87	17	
8.	do.	2	410	83		95	20	
9.	do.	2	400	86		87	18	

Table VII gives the effect of this volatile antiseptic on oats (first crop) and corn (second crop). The former showed an increase in growth in the treated soil; the latter was not affected.

TABLE VII.—Effect of carbon bisulphid on the growth of oats and corn

No.	Soil.	Carbon bisulphid added.	Weight of first crop, oats.			Weight of second crop, corn.		
			Green.	Dry.	Average.	Green.	Dry.	Average.
1.	Miami.	Per cent.	Gm.	Gm.	Gm.	Gm.	Gm.	Gm.
2.	do.	Control.	172	46.5	48.3	166	40	36
3.	do.	Control.	184	51		132	31	
4.	do.	Control.	171	46.7		118	29	
5.	do.	Control.	182	49		180	45	
6.	do.	2	200	59	57.8	155	37	37
7.	do.	2	205	59		161	38	
8.	do.	2	197	57.7		152	37	
9.	do.	2	192	57.5		135	30	

A general consideration of the data shows that corn in this soil type is apparently indifferent toward carbon bisulphid. Buckwheat, oats, and mustard were all benefited by the antiseptic.

#### BUCKWHEAT, MUSTARD, OATS, AND CORN IN DIFFERENT SOILS

The experiment with buckwheat, mustard, corn, and oats was a combination study of the effect of carbon bisulphid on bacterial activity and plant growth in three different soils. The first series contained Miami silt loam, the second series Miami soil diluted one-half by volume with sand, and the third series sand alone. According to chemical analysis, Miami silt loam is fairly rich in organic matter, nitrogen, potassium, and phosphorus. Of the three fertilizing elements, phosphorus perhaps is present in the smallest amount. The quantity of soil and its treatment was similar to that of the preceding experiment except that the treated jars were kept tightly covered with parchment paper. One month after the carbon bisulphid was added, these were removed. By

this means it was hoped to prevent a rapid volatilization of the anti-septic. The jars were not planted until three months after treatment.

At the beginning and at intervals of one, two, and three months bacterial activity was measured. Naturally, under the conditions of this experiment, carbon bisulphid proved very drastic. A great reduction in the number of bacteria, without any increase until the second month, was noted. The relation of carbon bisulphid to the number of bacteria was about the same in all three series. In the more compact type, Miami silt-loam soil, the carbon bisulphid proved most injurious to numbers, and consequently the period of increase was much later. Of the three soils, the treated sand showed the greatest proportional gain in number of bacteria.

Because of the severe nature of the carbon-bisulphid treatment, it was thought that probably the protozoa would be destroyed or the number greatly diminished. This was not the case, however, as protozoa were found in great numbers in both the treated and untreated soil.

Three months after treatment the jars were divided into two series and planted. The weights of the first and second crops are given in Tables VIII and IX.

TABLE VIII.—*Effect of carbon bisulphid on the growth of buckwheat and mustard in different types of soil*

No.	Soil.	Carbon bisulphid added.	Weight of first crop, buckwheat.			Weight of second crop, mustard.		
			Green.	Dry.	Average.	Green.	Dry.	Average.
1	Miami silt loam.....	Per cent.	Gm.	Gm.	Gm.	Gm.	Gm.	Gm.
2	.....do.....	Control.	123	22.5	21.5	12.0	3.4	3.3
3	.....do.....	Control.	107	20.5		10.0	3.3	
4	.....do.....	2	110	25.0	24.0	24.5	5.2	7.3
5	.....do.....	2	114	23.0		44.5	9.5	
	Half Miami silt loam, half sand.	Control.	74	15.5	15.25	24.0	3.75	3.67
6	.....do.....	Control.	72	15.0		17.0	3.60	
7	.....do.....	2	76	18.0	17.0	21.5	4.5	4.25
8	.....do.....	2	78	16.0		17.0	4.0	
9	Sand.....	Control.	20	2.5	2.75	4.0	0.4	.45
10	.....do.....	Control.	21	3.0		4.5	0.5	
11	.....do.....	2	21.5	3.0	3.00	17.5	1.2	.90
12	.....do.....	2	21.5	3.0		5.5	.6	

The figures of the buckwheat crop show the same general increase as noted in a previous experiment. Although not great, the gain in the treated series is consistent in all three soils.

The residual crop of mustard responded to a very marked degree to the carbon bisulphid treatment. In Miami silt loam the yield from the treated soil exceeded that of the control by more than 100 per cent. The gain in weight of oats in the treated soils was not so great, while the second-crop corn showed a loss (Table IX).



TABLE IX.—*Effect of carbon bisulphid on the growth of oats and corn in different types of soil*

No.	Soil.	Carbon bisulphid added.	Weight of first crop, oats.			Weight of second crop, corn.		
			Green.	Dry.	Average.	Green.	Dry.	Average.
1	Miami silt loam.....	Per cent.	Gm.	Gm.	Gm.	Gm.	Gm.	Gm.
2	.....do.....	Control.	162	47	47.7	114	25	24
3	.....do.....	Control.	180	48.5		103	23	
4	.....do.....	2	157	45.5	49	77	18.5	19.2
5	.....do.....	2	190	52.5		87	20	
6	Half Miami silt loam, half sand.....	Control.	82	26.5	27.2	76	16	15
7	.....do.....	Control.	85	28		56	14	
8	.....do.....	2	85	27.5	28	52	13	14
9	.....do.....	2	82	28.5		64	15	
10	Sand.....	Control.	18	6	6	12	4	4
11	.....do.....	Control.	18	5.8		14	5	
12	.....do.....	2	15	5.2	5.5	13.5	4	4.5

The results of the nitrate determinations agree with those obtained in previous experiments. At the time of planting the carbon-bisulphid soils were lower in nitrate but higher in ammonia than the original soil.

The data from Tables VIII and IX show that carbon bisulphid has a much more beneficial effect on mustard than on any other crop. Buckwheat and oats are benefited, but not so markedly as mustard. Corn fails to show any improvement from treatment with carbon bisulphid.

#### EFFECT OF CARBON BISULPHID ON BUCKWHEAT AND RAPE IN VARIOUS SOILS

The five soil types selected for the study of the effect of carbon bisulphid on buckwheat and rape in various soils ranged all the way from a very compact red clay to an open, sandy soil. After treating with 2 per cent of carbon bisulphid the soils were allowed to stand for three months before planting. Bacteria counts and nitrate determinations were made at the beginning and after two and three months. The effect of the carbon bisulphid on the total number of bacteria is very evident. In every case the carbon-bisulphid soil contained the most bacteria. The maximum gain occurred in the clay-loam soil, the minimum in the Norfolk sand. The increase due to the treatment was greatest after two months.

Here, again, the treated soils gave a much lower nitrate content than the controls. It seems safe to say that a rapid increase in numbers of bacteria in a carbon-bisulphid soil is followed by a decrease in the amount of nitrates.

Three months after treatment the soils were planted to buckwheat. Growth was slow at first, especially in the carbon-bisulphid series. The crop was harvested when 60 days old. The results of this experiment are shown in Table X.

TABLE X.—Effect of carbon bisulphid on the growth of buckwheat in different types of soil

No.	Soil.	Carbon bisulphid added.	Weight of first crop.		
			Green.	Dry.	Average.
1	Cecil clay	Per cent.	Gm.	Gm.	Gm.
2	do.	Control.	5	1.2	1.35
3	do.	Control.	7	1.5	
4	do.	2	12	3	3.5
5	do.	2	13	4	
6	Porters clay	Control.	10.5	2	2.75
7	do.	Control.	14.5	3.5	
8	do.	2	14.5	3.7	3.15
9	do.	2	10	2.7	
10	Clay loam	Control.	5.5	1.2	1.6
11	do.	Control.	12.5	2	
12	do.	2	28	6.5	7.25
13	do.	2	30.5	8	
14	Hagerstown loam	Control.	27	6.2	5.6
15	do.	Control.	25	5	
16	do.	2	17.5	4.5	6
17	do.	2	40.5	7.5	
18	Norfolk sand	Control.	32	8.7	9.45
19	do.	Control.	49.5	10.2	
20	do.	2	12	3	3.25
	do.	2	17.5	3.5	

With one exception, Norfolk sandy soil, the carbon-bisulphid series gave a larger yield. This was most marked in the case of clay-loam soil. The data on plant growth agreed with the plate counts.

The buckwheat was followed by a crop of Dwarf Essex rape. Unfortunately the young rape plants suffered seriously from insects. Although the tissue was too badly infested to save, a decided difference in growth could be seen. The beneficial effect of carbon bisulphid on rape was noted in every soil type.

#### EFFECT OF CARBON BISULPHID ON VARIOUS CROPS IN ACID SOILS

In order to study the effect of carbon bisulphid on the growth of higher plants in acid soils, a series of experiments was made. Four types of soil were selected for this work: Miami silt loam, Sparta sand, Colby silt loam, and Marshfield peat. The neutral Miami silt loam was used as a check for the acid soils. According to the Truog acidity test, Sparta sand requires 0.5227 gm. of calcium carbonate per 100 gm. of soil, Colby silt loam 1.021 gm., and Marshfield peat 4.43 gm. Four weeks after treatment with carbon bisulphid, the soils were planted.

#### RED CLOVER

The effect of carbon bisulphid on medium red clover in acid soils is clearly seen from the figures of Table XI. The clover grew luxuriantly in all soils except the untreated acid peat. Two crops were cut. Carbon bisulphid in peat soil caused an enormous gain in the growth of clover. This was very striking in both the first and second crop.

TABLE XI.—Effect of carbon bisulphid on the growth of red clover in acid soils

No.	Soil.	Carbon bisulphid added.	Weight of first crop, clover.			Weight of second crop, clover.		
			Green.	Dry.	Average green.	Green.	Dry.	Average.
		Per cent.	Gm.	Gm.	Gm.	Gm.	Gm.	Gm.
1	Miami silt loam.	Control.	138	(a)	139	129	19	20
2	do.	Control.	140			145	21	
3	do.	2	158			168	26	
4	do.	2	124	(a)	141	131	20	23
5	Sparta sand.	Control.	36			58	13	
6	do.	Control.	33			48	10	
7	do.	2	19	(a)	25	18	4	6
8	do.	2	31			43	8	
9	Colby silt.	Control.	95			110	20	
10	do.	Control.	87	(a)	91	85	14	17
11	do.	2	153			108	15	
12	do.	2	133			82	12	
13	Peat.	Control.	4	(a)	3	6	2.8	2.4
14	do.	Control.	2			5	2	
15	do.	2	83			53	9	
16	do.	2	79	(a)	81	46	8.5	8.7

(a) Lost.

Plate II, figure 2, shows the relative growth of clover in the treated and untreated soils.

Each figure for Miami silt loam in Table XI represents the average of triplicate jars. Because of the individual variation, it was decided to use 12 jars for this experiment. Six of these were used as controls and six treated with 2 per cent of carbon bisulphid. It is evident from the data that medium red clover in Miami soil is benefited both in the first and second crop by the antiseptic. In the Sparta sand a decrease was noted with each crop. The Colby silt loam gave a decided increase with the first crop, but not with the second.

Previous tests with these soils showed that the clover bacteria were present in sufficient numbers to produce good inoculation. In view of the large amount of carbon bisulphid applied, it was thought that this substance would probably injure nodule formation. However, examination of the root systems showed this was not the case. The plant roots were thoroughly inoculated, both in the treated and untreated soils. Apparently the plants in carbon bisulphid soils contained the greater number of nodules.

Because of the remarkable action of carbon bisulphid in peat soil, this part of the previous test was repeated. In addition to carbon bisulphid, the effect of flowers of sulphur was studied. If the data in the previous experiment are correct, the carbon bisulphid should greatly increase the growth of clover. A glance at the results in Table XII confirms this statement.

TABLE XII.—*Effect of carbon bisulphid and sulphur on the growth of red clover in peat soil*

No.	Soil.	Treatment.	Weight.		
			Green.	Dry.	Average.
			Gm.	Gm.	Gm.
1.	Peat.	Control	34	9	8.6
2.	do.	do.	30	8.2	
3.	do.	1 per cent of carbon bisulphid.	95	21.5	
4.	do.	do.	110	22	21.7
5.	do.	2 per cent of carbon bisulphid.	105	23	
6.	do.	do.	90	19.5	21.2
7.	do.	0.3 per cent of sulphur.	8	3.5	
8.	do.	do.	4	1	2.2

Carbon bisulphid causes a remarkable increase in the growth of clover on peat soil. There is apparently no decided difference in the action of 1 or 2 per cent of carbon bisulphid. Just why the volatile antiseptic should stimulate so markedly the growth of clover in the peat soil is not known. A more detailed study of the action of carbon bisulphid in peat is now under way. Flowers of sulphur at the rate of 0.3 per cent proved very injurious. In view of the high sulphur content of carbon bisulphid, it was thought that possibly free sulphur in peat might have somewhat the same effect.

## CORN AND MUSTARD

The action of carbon bisulphid on corn and mustard in acid soils was studied in an experiment the results of which are given in Table XIII.

TABLE XIII.—*Effect of carbon bisulphid on the growth of corn and mustard in acid soils*

No.	Soil.	Carbon bisulphid added.	Weight of corn.			Weight of mustard.		
			Green.	Dry.	Average.	Green.	Dry.	Average.
			Gm.	Gm.	Gm.	Gm.	Gm.	Gm.
1	Miami silt loam	Per cent.						
2	do.	Control.	190	50	64.5	83	18	19.5
3	do.	do.	360	79		145	21	
4	do.	2	315	79	65	159	27	25.5
5	do.	2	320	60		150	24	
6	Sparta sand	Control.	65	17	17.5	13	2.5	3
7	do.	Control.	70	18		19	5.5	
8	do.	2	100	21	23.5	11	2	2.5
9	do.	2	120	26		12.5	3	
10	Colby silt	Control.		83	84.5			10
11	do.	Control.	393	80		67	10	
12	do.	2	390	85	81	62	9.6	9.6
13	do.	2	385	77				
14	Peat	Control.	160	28	24	0		
15	do.	Control.	130	20		0		
16	do.	2	105	25	22.5	0		
17	do.	2		20		0		

It is clear from the data that carbon bisulphid does not materially benefit corn. An exception to this was seen in the case of Sparta sand; in this instance the treated series showed a slight improvement.

A comparison of the growth of mustard in acid and in neutral soil shows that this crop grows best in a neutral soil. In Sparta sand and Colby silt loam the yield of mustard in the treated soil was below that of the control, while in the peat soil it failed entirely. It seems very probable that the acid reaction of the soil inhibits the growth of mustard. For instance, Kossovich (7) reports that mustard is sensitive to acidity. The addition of 2 per cent of carbon bisulphid to Miami soil stimulated the growth of mustard. This agrees with the results of previous tests. An increase in the growth of mustard has been noted in all four experiments with carbon bisulphid in Miami soil.

One series of jars, corn on Miami silt loam, was replanted to buckwheat. As previously reported, buckwheat showed a distinct improvement in the carbon-bisulphid soil. If the control weights are taken as 100, the treated series is equal to 115.

A review of all the data on the effect of carbon bisulphid on higher plants shows very clearly that carbon bisulphid does not produce the same effect on all plants. In almost every case (except acid soils) the carbon bisulphid favors in a decisive way the growth of mustard. Next in order of their response to carbon bisulphid come rape, red clover, buckwheat, oats, and corn. In acid soils, especially those rich in organic matter, the growth of clover is greatly favored by the carbon-bisulphid treatment.

The majority of the evidence indicates that carbon bisulphid is most beneficial to the growth of higher plants in peat or in open, sandy soils.

#### EFFECT OF CARBON BISULPHID ON THE GROWTH OF PLANTS IN SILICA SAND

If carbon bisulphid is a plant stimulant, then the addition of the proper amount to a nutrient solution for plants should exert a beneficial effect on the growth of higher plants. To test this a series of experiments was performed on different plants.

##### BUCKWHEAT AND OATS

Eight jars were filled with pure silica sand (99 per cent pure quartz), and the following ingredients added to each jar:

Water ( $H_2O$ )	500	c.c.
Potassium nitrate ( $KNO_3$ )	5	gm.
Ferrous phosphate ( $Fe_3(PO_4)_2$ )	1.25	gm.
Calcium phosphate ( $Ca_3(PO_4)_2$ )	1.25	gm.
Calcium sulphate ( $CaSO_4$ )	1.25	gm.
Magnesium sulphate ( $MgSO_4$ )	1.25	gm.

In addition to the soluble plant food, half of the jars received 2 per cent of carbon bisulphid. After treatment the jars were held for two months before planting to buckwheat and oats. The results of the test are given in Table XIV.

TABLE XIV.—*Effect of carbon bisulphid on the growth of buckwheat and oats in silica sand*

No.	Carbon bisulphid added.	Weight of buckwheat.			Weight of oats.		
		Green.	Dry.	Average.	Green.	Dry.	Average.
	<i>Per cent.</i>	<i>Gm.</i>	<i>Gm.</i>	<i>Gm.</i>	<i>Gm.</i>	<i>Gm.</i>	<i>Gm.</i>
1.....	Control.	15.8	3.2	2.3	3.5	1.5	1.3
2.....	Control.	8.5	1.4		3.4	1.2	
3.....	2	37.5	7	5.6	.....	.....	6.5
4.....	2	21	4.2		21	6.5	

It is apparent from the data that carbon bisulphid in silica sand exerts a beneficial effect on the growth of both buckwheat and oats. This agrees with the results of Koch (6)—that carbon bisulphid stimulates the higher plant growth. Although the duplicate jars do not agree very closely, the highest yield of the control was lower than any of the treated groups. For some unexplainable reason, the oats in jar 3 failed to grow. The young seedling died soon after germination. Plate II, figure 3, is a reproduction of a photograph of the buckwheat series.

#### CLOVER, BUCKWHEAT, AND MUSTARD

The foregoing experiment was repeated, using 3-kgm. jars and Tollen's medium. Only 1 per cent of carbon bisulphid was added. The jars were planted 30 days after treatment. The yields of the different crops are presented in Table XV. From the beginning clover and mustard began to show the favorable effect of carbon bisulphid.

TABLE XV.—*Effect of carbon bisulphid on the growth of buckwheat, clover, and mustard in silica sand*

No.	Carbon bisulphid added.	Weight of buckwheat.			Weight of clover.			Weight of mustard.		
		Green.	Dry.	Average.	Green.	Dry.	Average.	Green.	Dry.	Average.
	<i>Per cent.</i>	<i>Gm.</i>	<i>Gm.</i>	<i>Gm.</i>	<i>Gm.</i>	<i>Gm.</i>	<i>Gm.</i>	<i>Gm.</i>	<i>Gm.</i>	<i>Gm.</i>
1	Control.	49	7.5	7	4.5	1	1.4	77	9	9
2	Control.	41	6.6		10	1.8		.....	.....	
3	1	49	7.8	7.6	12	2.2	2.2	52	5.8	8.6
4	1	45	7.5		12	2.3		98	11.5	

As compared with the results shown in Table XIV, the increase in the growth of buckwheat with carbon bisulphid was much smaller. The clover crop was about doubled in the presence of carbon bisulphid. Mus-

tard did not do well in sand cultures; growth was very irregular. Because of the size of the jars and the irregular growth of the crops it will be necessary to repeat the experiment.

#### EFFECT OF CARBON BISULPHID IN REINOCULATED SOIL

In the first part of this paper it has been shown that if soil treated with carbon bisulphid is reinoculated with fresh soil the bacterial processes are altered. The increase in number of bacteria attains a maximum much sooner and begins to decline earlier than in soil treated with carbon bisulphid but not reinoculated. This is also noted in the formation of soluble nitrogen. In order to record the effect on plant growth, the following experiment was planned. Six jars with 9 kgm. each of Miami silt-loam soil were used. Two months after treatment with carbon bisulphid, 2 per cent of untreated soil were added to jars 5 and 6. An equal amount was removed before the original soil was added. All of the jars were kept for another month before planting.

Plate counts three months from the date of treatment showed a decided increase in number of bacteria in the carbon-bisulphid soils. No appreciable difference existed between the carbon bisulphid and the carbon-bisulphid reinoculated soil.

The effect of treatment on nitrate content is evident from the following figures: If the nitrate nitrogen at the beginning is 100, then the control after three months is 370, carbon bisulphid is 50, and carbon bisulphid plus 2 per cent of the original soil is 44. Here, again, the inverse relation of number of bacteria and nitrate content is noted.

Protozoa were found in all of the soils and apparently in about the same number two months after treatment as in the original soil.

The effect of this treatment on the growth of oats and corn may be seen from the figures in Table XVI.

TABLE XVI.—*Effect of carbon bisulphid on the growth of oats and corn in reinoculated soil*

No.	Soil.	Treatment.	Weight of first crop, oats.			Weight of second crop, corn.		
			Green.	Dry.	Average.	Green.	Dry.	Average.
1	Miami	Control.....	Gm. 168	51	Gm. 50.9	Gm. 107	28	Gm. 29
2	do	do.....	178	50.75	52	108	30	29.5
3	do	2 per cent of carbon bisulphid.	178	50		86	33	
4	do	do.....	185	54	56.5	83	26	24
5	do	2 per cent carbon bisulphid plus 2 per cent of the original soil.	178	51.5		79	22	
6	do	do.....	215	61.5		101	26	

The average dry weight of oats in soil treated with carbon bisulphid was slightly greater than that of the control. This difference was most noticeable in the case of reinoculated soil. It appears that the reinoculation benefits the action of carbon bisulphid on the growth of oats. The second crop of corn gave the opposite results. The corn in untreated soil gave the highest yield.

#### EFFECT OF CARBON BISULPHID ON THE ACCUMULATION OF SULPHATES IN SOIL

Very soon after the jars were planted it was observed that the surface of carbon-bisulphid soil was partly covered with needle-like crystals. Qualitative tests showed that these were made up largely of sulphates, possibly magnesium sulphate. The occurrence of salts was noted in several of the soils treated with carbon bisulphid. Possibly a part of the carbon bisulphid was oxidized to sulphates. It has been reported that a small portion of the carbon bisulphid may be converted into sulphates (4, p. 247-251; 10, p. 151-152).

Samples of the treated and untreated soils were analyzed for sulphates.<sup>1</sup> The results are shown in Table XVII.

TABLE XVII.—Effect of carbon bisulphid on the accumulation of sulphates in the soil

No.	Time.	Treatment.	Sulphur as sulphates.
	<i>Months.</i>		<i>Per cent.</i>
1.	1	Untreated.	0.023
2.	1	2 per cent of carbon bisulphid.	.038
3.	3	Untreated.	.018
4.	3	2 per cent of carbon bisulphid.	.039
5.	5	Untreated.	.010
6.	5	2 per cent of carbon bisulphid.	.060

It is apparent from the data in this table that the addition of carbon bisulphid tends to increase the sulphate content of the soil.

#### CONCLUSIONS

The addition of carbon bisulphid to soil exerts a decided effect on the fauna and flora of the soil. This is characterized by a temporary reduction in the number of micro-organisms. Later, an enormous multiplication of bacteria takes place and an almost parallel increase in production of by-products or soluble nitrogen is noted. The ammonia content seems to follow the curve of bacterial growth and later gives way to larger amounts of nitrate. From the evidence it seems that carbon bisulphid in soil produces an increase in soluble compounds of nitrogen and sulphur.

<sup>1</sup> The author is indebted to Prof. W. E. Totttingham, of the Department of Agricultural Chemistry, for the analyses.



In Miami soil carbon bisulphid benefited the growth of buckwheat, oats, and mustard. No relation seems to exist between plant stimulation with carbon bisulphid and the form of the soluble nitrogen. In non-acid soils carbon bisulphid is most beneficial to sulphur crops. Mustard offers a good example. In all of the experiments, except acid soils, mustard showed an increased growth from the use of carbon bisulphid. Carbon bisulphid in peat soil greatly benefits the growth of red clover. In sand cultures plus soluble plant food carbon bisulphid favors the growth of certain plants.

The data show clearly that carbon bisulphid does not act alike in all soils or toward all crops.

## LITERATURE CITED

- (1) FRED, E. B.  
1911. Über die Beschleunigung der Lebenstätigkeit höherer und niederer Pflanzen durch kleine Giftnengen. *In* Centbl. Bakt. [etc.], Abt. 2, Bd. 31, No. 5/10, p. 185-245, 4 fig. *Literatur*, p. 242-245.
- (2) GAINNEY, P. L.  
1914. Effect of  $CS_2$  and toluol upon nitrification. *In* Centbl. Bakt. [etc.], Abt. 2, Bd. 39, No. 23/25, p. 584-595, 2 fig. *Literature*, p. 595.
- (3) GERLACH and VOGEL.  
1905. Ammoniakstickstoff als Pflanzennährstoff. *In* Centbl. Bakt. [etc.], Abt. 2, Bd. 14, No. 3/4, p. 124-128, 2 fig.
- (4) HEINZE, B.  
1907. Einige weitere Mitteilungen über den Schwefelkohlenstoff und die  $CS_2$ -Behandlung des Bodens. *In* Centbl. Bakt. [etc.], Abt. 2, Bd. 18, No. 1/3, p. 56-74, 1 fig.; No. 7/9, p. 246-264, 2 fig.; No. 13/15, p. 462-470; No. 19/21, p. 624-634; No. 24/25, p. 790-798.
- (5) HILTNER, LORENZ, and STÖRMER, KURT.  
1903. Studien über die Bakterienflora des Ackerbodens, mit besonderer Berücksichtigung ihres Verhaltens nach einer Behandlung mit Schwefelkohlenstoff und nach Brache. *In* Arb. Biol. Abt. Land- und Forstw. K. Gsndtsamt., Bd. 3, Heft 5, p. 445-545, 4 fig., pl. 9-10.
- (6) KOCH, ALFRED.  
1899. Untersuchungen über die Ursachen der Rebenmüdigkeit mit besonderer Berücksichtigung der Schwefelkohlenstoffbehandlung. *Arb. Deut. Landw. Gesell.*, Heft 40, 44 p., 5 pl.
- (7) KOSSOVICH, P. S., and ALTHAUSEN, L.  
1909. Influence of  $CaCO_3$  and  $MgCO_3$  on the soil and plants. (Abstract). *In* Exp. Sta. Rec., v. 23, no. 3, p. 226. 1910. Original article appeared in Trudui Mendelyevsk. Syezda Obsheh. i Prikl. Khim., god 1, 1907, p. 490-493. 1909. Not seen.
- (8) KRÜGER, WILHELM.  
1905. Über die Bedeutung der Nitrifikation für die Kulturpflanzen. *In* Landw. Jahrb., Bd. 34, Heft 5, p. 761-782, pl. 13-15.
- (9) LEHMANN, JUL.  
1875. Über die zur Ernährung der Pflanzen geeignetste Form des Stickstoffes. *In* Centbl. Agr. Chem., Bd. 7, p. 403-409.
- (10) MORITZ, J., and SCHERPE, R.  
1904. Über die Bodenbehandlung mit Schwefelkohlenstoff und ihre Einwirkung auf das Pflanzenwachstum. *In* Arb. Biol. Abt. Land- u. Forstw. K. Gsndtsamt., Bd. 4, Heft 2, p. 123-156.

- 
- (11) RUSSELL, E. J., and HUTCHINSON, H. B.  
1909. The effect of partial sterilisation of soil on the production of plant food.  
*In Jour. Agr. Sci.*, v. 3, pt. 2, p. 111-144, 4 fig., pl. 8-9.
- (12) ————  
1913. The effect of partial sterilisation of soil on the production of plant food.  
II. The limitation of bacterial numbers in normal soils and its consequences. *In Jour. Agr. Sci.*, v. 5, no. 2, p. 152-221, 7 fig.
- (13) SHERMAN, J. M.  
1914. The number and growth of protozoa in soil. *In Centbl. Bakt. [etc.]*,  
Abt. 2, Bd. 41, No. 18/23, p. 625-630.

# PLATE I

Plate cultures of soil organisms growing on agar:


Fig. 1.—Colonies of organisms from untreated soil.

Fig. 2.—Colonies from soil treated with 2 per cent of carbon bisulphid.

Fig. 3.—Colonies from soil treated with 2 per cent of carbon bisulphid and evaporated.

Fig. 4.—Colonies from soil treated with 2 per cent of carbon bisulphid, evaporated, and reinoculated with 5 per cent of soil from an untreated jar.





## PLATE II

Fig. 1.—Effect of varying amounts of carbon bisulphid on mustard; *A, B*, soil untreated; *C, D*, soil treated with 0.5 per cent of carbon bisulphid; *E, F*, soil treated with 1 per cent of carbon bisulphid; *G, H*, soil treated with 2 per cent of carbon bisulphid.

Fig. 2.—Effect of carbon bisulphid on clover in peat soil; *A, B*, soil untreated; *C, D*, soil treated with 2 per cent of carbon bisulphid.

Fig. 3.—Effect of carbon bisulphid on buckwheat in sand cultures; *A, B*, soil untreated; *C, D*, soil treated with 2 per cent of carbon bisulphid.



## CLIMATIC CONDITIONS AS RELATED TO *CERCOSPORA* *BETICOLA*<sup>1</sup>

By VENUS W. POOL, *Assistant Pathologist*, and M. B. MCKAY, *Scientific Assistant*,  
*Cotton and Truck Disease Investigations, Bureau of Plant Industry*

### INTRODUCTION<sup>2</sup>

Climatic conditions of both winter and summer bear an important relation to the vitality and development of *Cercospora beticola*. During cold weather certain conditions enable the fungus to overwinter, while certain other conditions are inimical to its growth, a fact which has an important bearing on the control of the disease, as the earliest infections on growing sugar beets (*Beta vulgaris*) originate from the overwintered fungus. In the early summer, after infection occurs, temperature, relative humidity, rainfall, and wind directly affect the development of the fungus, the rapidity of conidial production, and subsequent infection.

### OVERWINTERING

From the investigations here described it seems evident that under ordinary field conditions of winter the conidia of *C. beticola* usually live but a short time, although under ordinary herbarium conditions desiccation takes place only after exposure for several months. The sclerotia-like bodies (fig. 1, A, a), or masses of mycelium, the most resistant part of the fungus, which are embedded in the infected areas of the leaf blades and petioles, however, live over the winter under favorable conditions and in the spring produce conidia from the remnants of the old conidiophores (fig. 1, A, b), or both conidiophores and conidia (fig. 1, A, c) may be formed anew. For the purpose of making direct microscopical observation of such development sections of infected tissue which had been stored throughout the winter under favorable conditions were placed in hanging-drop cultures of bean agar. New conidiophores (fig. 1, B, b) grew from the masses of embedded mycelium, and although somewhat abnormal they produced rather typical conidia (fig. 1, B, c), thus showing that such material may be a source of early infection of growing plants.

<sup>1</sup> The investigations were carried on entirely in the field. Preliminary work was conducted during 1911 and 1912 at Rocky Ford, Colo. The detailed data were collected during 1912 and 1913 at Rocky Ford, which is in the Arkansas Valley of Colorado, a semiarid region under irrigation, and during 1914 near Madison, Wis., where the rainfall and average humidity were greater.

<sup>2</sup> The writers are indebted to Mrs. Nellie E. Fealy, of the Bureau of Plant Industry, for aid in editing and revising the manuscript.






FIG. 1.—*Cercospora heliicola*: A, Section of overwintered sugar-beet leaf showing embedded sclerotia-like body, a, with a mass of old conidiophores, b, from which a new conidium, c, was produced. B, Production of rather typical conidiophores, b, and conidia, c, from a sclerotia-like mass, a, taken from overwintered sugar-beet leaf.

## CONIDIA

Thümen (1886, p. 50-54)<sup>1</sup> believed that the spores of *Cercospora beticola* are able to live for a certain length of time in the soil and retain their viability and produce new infection, and Pammel (1891, p. 238-243) and Massee (1906, p. 52-53) accord with this view. In the investigations here considered it was found that when kept dry, as in the case of herbarium material, the conidia remained viable for 8 months (Table I, tests 10 to 13), but soon after that no growth occurred. Only rarely were conidia found on the infected areas of the leaves which were exposed to outdoor weather conditions, and such conidia seemed to lose their vitality soon after harvest. No germination was found to take place under optimum conditions in the case of conidia which had been thus exposed from 1 to 4 months (tests 14 and 15). However, conidia occasionally found on spots that had been well protected, for instance in the interior of a pile of hayed beet tops, retained their viability for from 5 to less than 12 months (tests 16 and 17). Since the conidia are rarely found after a short time even on infected material that has been well protected and since they rarely germinate after being exposed outdoors for even 1 month after harvest, it would seem that under ordinary field conditions they play no important part in the overwintering of the fungus.

TABLE I.—Viability of the conidia of *Cercospora beticola* as affected by desiccation

Test No.	Environment.	Period of exposure.	Viability.
1	Stored, dry.....	14 years.....	None.
2	do.....	11 years.....	Do.
3	do.....	10 years.....	Do.
4	do.....	5 years.....	Do.
5	do.....	4 years.....	Do.
6	do.....	3 years.....	Do.
7	do.....	2 years.....	Do.
8	do.....	11 months.....	Do.
9	do.....	10 months.....	Do.
10	do.....	8 months.....	Slightly viable.
11	do.....	7 months.....	Extremely viable.
12	do.....	6 months.....	Do.
13	do.....	5 months.....	Do.
14	Left in field after harvest, Wisconsin.....	1 month.....	None.
15	Left in field after harvest, Colorado.....	4 months.....	Do.
16	Stored inside pile of hayed sugar-beet leaves.....	5 months.....	Extremely viable.
17	do.....	12 months.....	None.

## SCLEROTIA AND MYCELIUM

Various investigators have attempted to determine whether different fungi live in the soil over winter and the manner in which they overwinter. Treboux (1914) found that the mycelia of several different rusts overwinter on host material freely exposed to climatic conditions. Stewart (1913) placed in boxes of soil potato leaves and tubers infected with *Phytophthora infestans*, exposed them to outdoor winter conditions, and found that plants grown on such soil developed no blight. However,

<sup>1</sup> Bibliographic citations in parentheses refer to "Literature cited," p. 60.

temperature and moisture conditions in boxes of soil exposed above-ground to winter conditions are much more varied than in soil at different depths in the field where normal overwintering usually occurs.

In the overwintering experiments here described the host material was kept in an environment comparable to ordinary field conditions. The experiments at Rocky Ford, Colo., were started about the middle of October, 1912, and continued for 11 months. In these experiments some of the infected material was mixed with soil, placed in boxes, and exposed above-ground during the winter (Pl. III, 1); a second portion was buried from 1 to 8 inches in the ground (Pl. III, 2), wire netting being used above and below the infected material to insure ready location when examinations were made for cultural tests (Pool and McKay, 1915); a third portion of the infected tops was placed in a pile on top of the ground (Pl. III, 3). During the experiment records were kept of soil and air temperatures, the former being taken at a depth of 5 inches and the latter being obtained from the Weather Bureau station at Rocky Ford.<sup>1</sup>

The experiments carried on near Madison, Wis., were started the last of November, 1913, and continued through the winter. Infected sugar-beet tops were buried in the soil at depths of 5 and 8 inches, while seed-beet stalks were left under ordinary conditions in the field. In this experiment also records were kept of soil and air temperatures, the former being taken from March until June at a depth of 5 inches and the latter obtained from the Weather Bureau station at Madison.

The effect of desiccation on material kept under herbarium conditions was to kill probably all life of the fungus within 12 months, as already shown, but material kept under an environment having more or less moisture accompanied by the disintegrating action of various organisms was affected in an entirely different manner, as will be shown. All cultures from the infected material used in the two experiments above outlined were made from definite leaf-spots. Although the diseased tissue was the last to be completely disorganized and consequently could be found as long as any portion of the leaf remained, it became more and more difficult to obtain such tissue as time went on.

The fungus was unable to survive six months' outdoor exposure in boxes of soil (Table II, experiment 2), and this was also true of the fungus on leaves which had been freely exposed to outdoor conditions—for instance, on the outside of a hayed pile of sugar-beet tops (experiment 3), and on leaves buried 6, 7, and 8 inches in the ground (experiments 19 to 23). In cultures from infected mother-beet stalks and leaves that had been left in the field for a time and then plowed under or stored there was no growth, or only an indefinite growth, of the fungus after 7 months (experiments 8 to 10), while in infected material that had been protected in the interior of a pile of hayed beet tops (experiment 4) and in material

<sup>1</sup> All the records included in this paper from the Weather Bureau station at Rocky Ford, Colo., were kindly furnished by Mr. F. K. Blinn, the local observer.

that had been slightly covered or buried from 1 to 5 inches in the ground the life of the fungus was entirely extinct after 12 months (experiments 11 to 18). The death of the fungus in material plowed under is due in all probability to the rapid disorganization which results under favorable temperature and moisture conditions, such, for instance, as those which prevailed at Rocky Ford through the winter of 1912-13. During that period there was insufficient moisture to permit severe freezing, but there was a daily extreme variation of soil temperature, indicating that the air temperature produced the changes through the more or less dry soil. In the experiments at Madison there was only a partial disintegration of the buried beet tops six months after harvest, but other factors impaired the vitality of the fungus and its life appeared to be entirely extinct; consequently, notwithstanding the great differences in soil factors, comparable results as to the life of the fungus were obtained from the experiments at both places.

TABLE II.—Effect of desiccation and overwintering on the viability of *Cercospora beticola* in infected sugar-beet tops under field conditions at Rocky Ford, Colo., and Madison, Wis.

Experiment No.	Environment of sugar-beet-top material.	Period of exposure.	Number of spots from which cultures were made.	Number of viable spots.	Condition of leaves.
1	Dried, stored:				
	Illinois, Iowa .....	14 years ..	10	0	Good.
	Connecticut .....	11 years ..	10	0	Do.
	New York .....	10 years ..	12	0	Do.
	Wisconsin .....	5 years ..	10	0	Do.
	Iowa .....	4 years ..	10	0	Do.
	Maryland .....	3 years ..	10	0	Do.
	Colorado .....	2 years ..	10	0	Do.
	New Jersey .....	10 months ..	10	3	Do.
	Colorado .....	9 to 11 months ..	20	10	Do.
2	Stored in soil in boxes and left free under outdoor conditions, Colorado.	2 months ..	15	15	Do.
		3 months ..	7	7	Do.
		4 months ..	13	4	Do.
		5 months ..	12	2	Do.
		5½ months ..	12	0	Do.
3	From the outside of "hayed" pile of sugar-beet tops, Colorado.	7 months ..	6	0	Do.
		10 months ..	13	0	Do.
		12 months ..	10	10	Do.
		3 months ..	66	64	Do.
		4 months ..	29	27	Do.
4	From the interior of "hayed" pile of sugar-beet tops, Colorado.	5 months ..	40	40	Do.
		7 months ..	18	12	Do.
		10 months ..	15	10	Do.
		12 months ..	25	0	Do.
		13 months ..	10	10	Do.
5	In field, Colorado .....	5 months ..	11	8	Do.
		8 months ..	10	2	Do.
6	Leaves from "mother beet" stalks free in field, Wisconsin.	5 months ..	21	15	Do.
7	First-year sugar-beet leaves free in field, Wisconsin.	5 months ..	32	3	Do.
8	Spots on "mother beet" stalks free in field, Wisconsin.	8 months ..	40	0	Partially disintegrated.
9	Spots on "mother beet" stalks free in field 6 months, then plowed under 1 month, Wisconsin.	4 months ..	7	0	Good.
		7 months ..	35	42	Do.
		7 months ..	10	0	Somewhat softened.

<sup>a</sup> Herbarium specimens for this test were furnished by Barrett, Illinois; Clinton, Connecticut; Whetzel, New York; Pammel, Iowa; Norton, Maryland; and Cook, New Jersey.

<sup>b</sup> 395 colonies.

TABLE II.—Effect of desiccation and overwintering on the viability of *Cercospora beticola* in infected sugar-beet tops under field conditions at Rocky Ford, Colo., and Madison, Wis.—Continued

Ex- per- iment No.	Environment of beet-top material.	Period of exposure.	Number of spots from which cultures were made.	Number of viable spots.	Condition of leaves.
10	Spots on "mother beet" stalks free in field 4 months, then stored dry 3 months, Wisconsin.	7 months..	10	5 <sup>2</sup>	Good.
11	Buried 2 inches in ground, Colorado.	6 months..	21	0	Partially disintegrated.
		7 months..	10	2	Do.
		10 months..	8	3	Greatly disintegrated.
12	Buried 2 inches in ground, Colorado.	10 months..	24	0	Entirely disintegrated.
		5 months..	14	2	Partially disintegrated.
		8 months..	19	10	Do.
13	Buried 3 inches in ground, Colorado.	10 months..	11	0	Greatly disintegrated.
		12 months..	28	0	Entirely disintegrated.
		16 months..	21	10	Partially disintegrated.
14	Buried 3 inches in ground, Colorado.	10 months..	11	1	Greatly disintegrated.
		12 months..	18	0	Entirely disintegrated.
		16 months..	14	0	Greatly disintegrated.
15	Broken and buried 3 inches in ground, Colorado.	7 months..	12	8	Do.
		10 months..	16	0	Entirely disintegrated.
		16 months..	19	5	Partially disintegrated.
16	Buried 4 inches in ground, Colorado.	10 months..	19	1	Greatly disintegrated.
		6 months..	20	4	Do.
		10 months..	15	0	Entirely disintegrated.
17	Buried 5 inches in ground, Colorado.	12 months..	20	0	Do.
		4 months..	30	0	Good.
		5 months..	35	37	Do.
18	Buried 5 inches in ground, Wisconsin.	6 months..	87	0	Do.
		7 months..	80	0	Partially disintegrated.
		15 months..	30	37	Do.
19	Plowed under in field, about 5 inches, Wisconsin.	15 months..	20	0	Do.
		5 months..	8	0	Greatly disintegrated.
		6 months..	20	4	Do.
20	Buried 6 inches in ground, Colorado.	10 months..	12	0	Do.
		12 months..	15	0	Do.
		16 months..	18	0	Do.
21	Broken and buried 6 inches in ground, Colorado.	7 months..	10	0	Do.
		10 months..	16	0	Entirely disintegrated.
		16 months..	15	0	Do.
22	Buried 7 inches in ground, Colorado.	7 months..	10	0	Do.
		10 months..	12	0	Do.
		12 months..	22	0	Do.
23	Buried 8 inches in ground, Colorado.	10 months..	20	0	Do.
		7 months..	10	0	Do.
		10 months..	12	0	Do.
24	Buried 8 inches in ground, Wisconsin.	4 months..	20	0	Good; ground frozen.
		5 months..	35	27	Good.
		6 months..	27	0	Good; leaves have sour odor.
		6 1/2 months	100	0	Partially disintegrated.

These experiments and observations made in the field during several spring and summer months showed that on leaves slightly protected on or near the surface of the ground during the winter *C. beticola* can live a sufficient length of time to be a source of infection for the succeeding sugar-beet crop and that the fungus is entirely killed by planting time when the infected material is plowed under to a depth of 6 to 8 inches in the fall.

#### AIR AND SOIL TEMPERATURES AT ROCKY FORD, COLO., AND AT MADISON, WIS.

Comparison of the air and soil temperatures which prevailed during the experiment at Rocky Ford and Madison showed a wide difference.

One of the most striking characteristics of these temperatures at Rocky Ford was the wide range between the maximum and the minimum, and this range may be observed throughout the entire records (fig. 2, 3). In the case of the soil temperatures especially, the wide range appeared to be due to a lack of moisture, the extreme variations being greater than if more moisture had been present. A comparison of the records shows that the variation in air temperature was much less and the mean daily temperature constantly lower at Madison than at Rocky Ford, notwithstanding the fact that the daily minimum temperature was usually lower at Rocky Ford. A comparison of the soil temperatures at the two points, however, shows that at Madison it probably remained more constant and was never as low as at Rocky Ford. This was due apparently to the greater amount of moisture in the soil at Madison and consequently its continued frozen condition. After March 23, the date on which the record was begun at Madison, the soil temperature at that place was never below 29° F., notwithstanding the fact that the air temperature was as low as 15° on April 8, while the minimum soil temperature at Rocky Ford was 22° on December 21 and 25° on February 8. However, as the air temperature on these dates was lower here than at Madison, comparisons can not be drawn too closely.

In view of the presence of snow on the ground, which, as is well known, protects the soil from the extreme variations of air temperature, and the prevailing low air temperatures, as shown by the records, it may be assumed that the soil temperatures at Madison during January and February and the early part of March varied but little from freezing. This assumption is supported by Frödin's experiments (1913), which showed in general that when the air temperature was much lower than that of the soil the soil temperature in ground covered with snow was higher than in bare ground. He found that temperatures taken at a depth of 10 cm. in the former were the same as those taken at a depth of 27.4 cm. in the latter. After the early part of April the minimum soil temperatures at Rocky Ford and Madison agreed closely, although the minimum air temperature at the former place remained generally the lowest of the temperatures recorded.

Temperatures obtained from the interior of a pile of hayed sugar-beet leaves by means of a soil thermograph buried in the pile varied less than temperatures taken outside the pile, as shown by the following records made on May 8, 1913, and as was probably the case during the entire winter season: Temperature inside pile, maximum, 67° F.; minimum, 58°; difference, 9°. Temperature outside pile, maximum, 84° F.; minimum, 45°; difference, 39°.

In view of the fact that the fungus lived twice as long inside the pile as it did on the outside it would seem that a more uniform temperature might be regarded as one of the controlling factors in the life of the fungus.






Fig. 3.—Curves of the maximum and minimum soil and air temperatures for Rocky Ford, Colo., from March 13 to June 17, 1931, and for Madison, Wis., from March 13 to June 17, 1934



Low temperatures are not entirely inhibitive, as was shown by thermal tests of artificial cultures. After such cultures had been exposed to temperatures averaging  $0.9^{\circ}$  C. for 48 days and then kept at  $28^{\circ}$  C., numerous colonies developed. Also, heavily infected leaves kept at  $0.9^{\circ}$  C. for 97 days yielded good growth when cultures were made and held at favorable temperatures. Had the cultures been exposed to freezing temperatures or to extreme variations in temperature, the effect would doubtless have been more pronounced.

Although the temperature variations and the amount of soil moisture at Rocky Ford and Madison differed greatly, the effect on the life of the fungus was apparently the same at both places. It may be concluded that conditions of the soil which favor the process of disintegration are the most important factors in the control of the disease, and these experiments indicate that these processes are most active at a depth of 6 to 8 inches.

#### SUMMER CLIMATIC CONDITIONS

The summer climatic conditions here considered were recorded during 1913 in fields of first-year sugar beets grown at Rocky Ford, these fields being an example of the usual progress of the disease where neither rotation nor sanitation at the preceding harvest time had been practiced.

A study of the temperature and humidity records taken at different places in a beet field at Rocky Ford and at the Weather Bureau station 3 miles from the field was made to determine their comparative values in making important correlations. The records made in the sugar-beet field were taken by means of hydrothermographs kept in meteorological instrument shelters 5 feet above the ground (Pl. IV, fig. 1) and among the plants (Pl. IV, fig. 2). These were checked at frequent intervals with a sling and cog psychrometer (Shaw, 1914), respectively, and under Colorado conditions were found to be accurate. The records of the Weather Bureau station were taken by means of maximum and minimum thermometers kept in an instrument case about 5 feet above the ground in an open space (fig. 3).

The daily maximum and minimum temperatures and humidities, together with the total number of hours the humidity was above 60 from noon of the preceding day to noon of the given day, are used in the present interpretations. It has been found that when a high relative humidity prevails, the stomata of the sugar-beet leaves are usually open; and as the fungus enters the leaves only through the open stomata, the length of time they remain open is a fundamental factor in determining the possible occurrence of infection (Pool and McKay, 1916).

#### AIR TEMPERATURE AND RELATIVE HUMIDITY

The temperature and relative humidity taken with hygrothermographs placed near the ground among the plants varied widely from those taken with hygrothermographs in the air above the field and also from those

taken at the Weather Bureau station; hence, the place where the records were taken for use in the present correlations with the development of the disease is an important consideration.

**AIR TEMPERATURE.**—At Rocky Ford the maximum temperatures taken among the plants near the surface of the ground from June 13 to 30 ranged from 2 to 19 degrees higher and the minimum temperatures generally from 1 to 14 degrees lower than those taken at 5 feet above the ground (fig. 4). This was due to the fact that the plants were small during this period and covered only a portion of the ground; consequently during the daytime the temperature of the soil became higher than that of the air, and in turn the temperature of the air near the ground became higher than that of the air a few feet above. During the night the reverse occurred, the surface soil losing its heat by radiation and conduction faster and finally reaching a lower temperature than that of the air in contact with it, after which the heat of the latter gradually passed into the soil and as a result the temperature of the air immediately above the ground eventually became lower than that a few feet higher up. It is possible that convection currents also tended to lower the temperature of the air immediately above the ground; for, as is well known, when it is not disturbed by other factors, the coolest air settles to the lowest levels.

The maximum temperatures of the air near the ground, as shown by the records, were higher for a longer period during June than at any time during the season, varying from 100° to 106° F. on nine different days between the 14th and 26th of that month and rising above 100° only once thereafter, on August 16. The maximum temperature of the air 5 feet above the ground, on the other hand, was lower during June than during the middle of the season, ranging from 90° to 93° on six different days during the month, while it was above 90° and sometimes as high as 100° on 12 different days during July.

As shown by the records, the temperature of the air near the ground among the plants was lower during the middle than during the early part of the season. This was probably due to the difference in the size of the plants, the larger plants practically covering the ground in mid-season and preventing the heating of the surface soil, while early in the season the smaller plants covered the ground but sparsely and consequently afforded less protection against heating. Comparison of the records also shows that during the middle of the season the temperature of the air among the plants near the ground was practically the same as that of the air 5 feet above the field and that throughout the entire period the latter was quite comparable with the temperatures taken at the Weather Bureau station (fig. 4).

A similar marked variation was shown at Madison, the maximum temperature there being almost constantly higher and the minimum tempera-



ture usually lower among the beet plants than the temperature shown by the Weather Bureau records, which were taken on top of a four-story building about a mile from the sugar-beet field. These wide variations between the air temperature taken near the ground among the plants and that taken 5 feet above the field and between the former and the temperature taken at the Weather Bureau stations show that for correlation with fungous activities only the records taken among the plants should be used.

RELATIVE HUMIDITY.—There was also a wide variation in the humidity near the ground among the plants and 5 feet above the field. For instance, the daily minimum humidity at Rocky Ford from June 13 to 29, with two exceptions, was higher and remained above 60 generally for a longer period in the air above the plants than among the leaves near the ground (fig. 4), owing to the higher temperature at the surface of the ground as a result of the small amount of covering afforded by the young plants. During this period the daily variation of humidity among the leaves was extreme, ranging from 99 to 10 on June 13, from 99 to 16 on June 25, and from 100 to 8 on July 2. After June 29, on the other hand, the minimum humidity was generally higher, the humidity remained above 60 for a longer time among the leaves than in the air above, and the daily variation among the leaves was less extreme than earlier in the season. These conditions were due mainly to the greater amount of covering afforded by the larger plants and consequent longer retention of moisture among the leaves. The humidity both among the plants and in the air 5 feet above the field remained, on an average, above 60 for a longer time each day during midsummer than during June, owing in part to the increased use of irrigation water as the season advanced and the increased amount of moisture in the surrounding air resulting from the increased transpiration of the larger plants.

Comparison of the Madison and the Rocky Ford records (fig. 5) of the number of hours that the relative humidity remained above 60 each day among the sugar-beet plants shows that throughout the season it was higher, on an average, at Madison. Here it remained above 60 for a longer time each day during the latter half of June, when the records were started, and for a shorter time each day during August than during any other summer month. This was due to difference in the amount of rainfall, there being frequent rains during the former period and comparatively dry weather during the latter. At Rocky Ford the facts were reversed, the humidity remaining above 60 for a longer time each day during midseason than during the latter half of June or the first part of September. This was probably due to more frequent irrigation and the increased covering afforded by the larger plants of midseason.

Table III shows the average number of hours a day that the relative humidity remained above 60 at Madison and at Rocky Ford.

TABLE III.—Average number of hours a day that the relative humidity was above 60 at Madison, Wis., and Rocky Ford, Colo., during the summer of 1914 and 1913, respectively

Date.	Madison, Wis. (1914).	Rocky Ford, Colo. (1913).
	Hours.	Hours.
June 16 to 30.....	19.4	16.8
July.....	17.3	14.2
August.....	16.6	14.1
September 1 to 6.....	18.6	11.3
Seasonal average.....	17.4	13.4




FIG. 5.—Curves of the maximum and minimum temperatures among sugar-beet plants and at the Weather Bureau station, and the seasonal rainfall records at Madison, Wis., in 1914, and the number of hours that the humidity remained above 60 among the sugar-beet plants in the field at Madison, Wis., in 1914, and at Rocky Ford, Colo., in 1913.

The greater average number of hours of high humidity at Madison accounts for the periods of extreme infection which occurred there when the fungus was present. Here leaves badly infected with *Cercospora beticola* and entirely covered with conidia were found at times, but this condition was rarely seen at Rocky Ford. There were numerous cases of cotyledon infections also at Madison, the high humidity early in the season favoring their occurrence; but no such infections were found at Rocky Ford.

## RAINFALL AND IRRIGATION

The rainfall records made during the summer season of 1913 in the beet field at Rocky Ford in which infection was studied in detail (fig. 4, 7) were obtained by means of a rain gauge placed at the edge of the sugar-beet field (Pl. IV, fig. 1). Most of the rain was in the form of local showers, the amount varying greatly within a radius of less than 2 miles; but occasionally general rains fell. The effect of the increased relative humidity resulting from rainfall usually lasted longer among the leaves than in the air 5 feet above (fig. 7).

The effect of irrigation on humidity was found to be similar to the effect of rain. On July 2, before the field was irrigated, its humidity was as low as 8 and on July 3 and 4 remained above 60 for 7 and 6 hours, respectively. On July 4 and 5 the field was irrigated and the humidity remained above 72 on the 4th and above 60 during 23 hours of the 5th. On July 27 the field was again irrigated and the humidity remained above 60 for 15 hours that day and 21 hours the following day. On August 19 and 20 the field was irrigated the third time and the humidity remained above 60 for 12 and 13 hours, respectively, and the next day 21 hours. The general humid conditions necessary for leafspot infection, however, are developed much better by rain than by irrigation, because of the latter being comparatively local and unaccompanied by the atmospheric conditions attending rainfall.

## WIND

Records of wind velocity at Rocky Ford were taken by means of an anemometer placed 6 feet in the air at the edge of the beet field (Pl. IV, fig. 1), the readings being made at irregular intervals and the velocities computed being the hourly averages from one reading to the next. As the records were not made daily, accurate hourly velocities for different intervals during the day can not be obtained from the records. They show, however, that the average seasonal velocity from June 12 to September 22 was 5.3 miles per hour. Occasionally two daily readings were made, one in the morning and the second late in the afternoon. These show that the average velocity of the wind was always higher during the day than at night, the greatest velocity usually prevailing in the afternoon during the period of lowest humidity. While no general dissemination of conidia was correlated with high wind velocity, the afternoon combination of highest wind with lowest humidity apparently favored the dissemination of conidia. In fact, in the case of air cultures made at different times during several days, it was found that the fungus grew usually only on those exposed during the afternoon.

## SUMMER INFECTION CYCLES

The thermal relations of the fungus are closely linked with the effect of various climatic factors on the production and dissemination of conidia and on infection cycles. With a view to determining these relations the fungus was grown in Petri-dish cultures in thermostats at different and varied temperatures. At first the moisture was probably more or less constant, but as time went on it became relatively low. The effect of different temperatures, however, was comparable with that observed under existing field conditions.

## THERMAL RELATIONS OF THE FUNGUS IN CULTURES

Tests of the fungus on string-bean agar were made at Washington during November and December, 1913, and January, 1914. The cultures were obtained from isolations made at the time of the tests from infected sugar-beet leaves collected at Rocky Ford during the preceding September. One colony of the first isolations was macerated in 10 c. c. of sterile water, and one platinum loop of this suspension was used for each tube of medium. Three poured plates were used for each single test. The cultures were exposed to different constant temperatures and to varied constant temperatures (high and low changed to low and high, respectively). Exposures were also made for 8 hours at the higher temperatures and then for 16 hours at lower temperatures, and, after a short interval of exposure in a certain number of these tests, both temperatures were lowered, it being possible in this way to approximate night and day temperatures in the field under normal conditions.

**SERIES A (DIFFERENT CONSTANT TEMPERATURES).**—When the cultures were held at different constant temperatures, the abundance and size of the individual colonies gradually increased, while the time necessary for development decreased with the temperatures 12.5°, 17.3°, 19.2°, 20°, and 30.8° C. The best growth was made at a temperature of 30.8°, but this in all probability was slightly above the optimum constant temperature, as no growth took place in cultures held for 9 days at 34.7°, 35.8°, and 40.6°, respectively (Table III, series A).

**SERIES B (VARIED CONSTANT TEMPERATURES).**—Although no growth of the fungus took place in cultures held at constant temperatures of 34.7° and 35.5° C., a small percentage of normal colonies developed in cultures exposed for three days to these temperatures and then for several days to a temperature of 30.8°, while in cultures exposed for three days to 40.5° no growth occurred when subsequently held at 30.8°. On the other hand, in cultures exposed for three days to a temperature of 30.8° there was almost a normal development of the colonies for three days after they were exposed to 34.7°, but at the end of five days the inhibitive effect of the latter temperature became manifest. In the case of cultures

changed from 30.8° to 40.5° only a very slight increase in growth was apparent during the first three days at the higher temperature, and after that it ceased entirely (Table III, series B).

TABLE III.—Comparative diameter of colony growth (in millimeters) of *Cercospora beticola* at different constant temperatures,<sup>a</sup> at decreasing and increasing constant temperatures, and at daily varied temperatures

SERIES A, CONSTANT TEMPERATURES												
Diameter of colony growth at temperature (°C.) of—												
Maximum.....	2.1	7.9	11.1	14.4	19.1	20.2	21.0	33.0	36.2	37.4	41.0	
Minimum.....	0.3	4.0	8.0	11.5	16.3	17.6	19.5	29.7	33.8	34.5	40.0	
Average.....	0.9	5.4	8.9	12.5	17.3	19.2	20.0	30.8	34.7	35.8	40.6	
Period of growth:												
3 days.....	0	0	0	0	0.25	0.65	0.67	3.6	0	0	0	
6 days.....	0	0	.25	.8	1.2	2.1	6.4	0	0	0	0	
9 days.....	0	0	.25	.6	1.5	5	4.6	7	0	0	0	
11 days.....	0	0	.4	.85	2.8	7	6.4					
14 days.....	0	0	.5	1.5	4.4							
18 days.....	0	0	.7	2.2								
22 days.....	0	0	1.2	3								

<sup>a</sup> The temperatures of each thermostat for all tests were averaged from two daily readings continued throughout the time of the experiment.

<sup>b</sup> No growth occurred in these plates when held at 28° C. for 10 days.

SERIES B, DECREASING AND INCREASING CONSTANT TEMPERATURES.

Diameter of colony growth at temperature (°C.) of—					
	<sup>a</sup> 34.7	<sup>a</sup> 35.5	<sup>a</sup> 40.5	<sup>b</sup> 30.8	<sup>c</sup> 30.8
Period of growth:					
3 days.....	0	0	0	3.6	3.4
6 days.....	4	3.6	0	0	3.7
8 days.....	d 8.3	e 9.3	0	6.4	3.7

<sup>a</sup> Temperature changed to 30.8° C. after three days.

<sup>b</sup> Temperature changed to 34.7° C. after three days.

<sup>c</sup> Temperature changed to 40.5° C. after three days.

<sup>d</sup> Only 19.2 per cent of the normal number of colonies developed.

<sup>e</sup> Only 12.8 per cent of the normal number of colonies developed.

SERIES C, DAILY VARIED TEMPERATURES

Diameter of colony growth at temperature (°C.) of—							
	14.5	14.5	14.5	20	20	20	20
16 hours at.....				30.8	34.7	35.8	40.6
8 hours at.....	19.2	21.6	28				
Period of growth:							
3 days.....	0.4	0.5	0.7	1.7	0.6	0.6	0
5 days.....	.7	1.9	3.4	6	3.6	3.2	0
7 days.....	2.7	3.6	5.2	8.8	4.8	5.4	0
9 days.....							0

SERIES C (DAILY VARIED TEMPERATURES).—In these tests the temperatures were made to correspond closely with summer outdoor temperatures of night and day by holding the cultures for 16 hours at the lower and for 8 hours at the higher. After seven days' exposure the growth of colonies



on cultures exposed to temperatures of 14.5° and 19.2° C. averaged 2.7 mm. in diameter; after exposure for the same length of time to 14.5° and 21.6°, 14.5° and 28°, and 14.5° and 30.8° the growth gradually increased until it reached a maximum diameter of 8.8 mm; but when exposed to higher temperatures (20° and 34.7° or 20° and 35.8°) the growth gradually diminished until finally it equaled approximately that attained under 14.5° and 28°. There was no growth on cultures exposed for nine days or longer to 20° and 40.6° (Table III, series C).

**SERIES D (HIGH VARIED CHANGED TO LOW VARIED TEMPERATURES).**—A plate culture exposed for three days to temperatures of 20° and 40.5° C., being held 16 hours at the lower and 8 hours at the higher, and then for six days at 20° and 30.6°, developed 23 colonies, averaging 10.3 mm. by the end of the latter period, while a check plate exposed constantly to a temperature of 30.6° developed 100 colonies by the end of the latter period. A plate exposed to the higher temperatures—20° and 40.5°—for five days and then held at 20° and 30.6° for six days developed six colonies at the end of the latter period, while a plate exposed to 20° and 40.5° and then held at 20° and 30.6° for seven days developed no growth of the fungus.

Later on in this paper the fact that high minimum and maximum temperatures inhibit the growth of the fungus, as brought out by these tests, is correlated with the effect of existing high field temperatures, with their consequent accompanying factors, on the leaf spot. Although the optimum temperature variations—20° and 30.8° C.—were found to be very favorable to the development of leafspot in the field, little or no increase in the disease was observed to follow high night and day field temperatures—20° and 40.5°, respectively.

It was also observed that different temperatures affect conidial septation. The normal average septation varies from 6 to 11, but during warm, humid periods the conidia were usually found to be many septate, sometimes as high as 20-septate, while after a cooler period, such as usually occurs in September, they were only from 2- to 4-septate.

#### RELATION OF CONIDIAL PRODUCTION AND DISSEMINATION TO CLIMATIC CONDITIONS

For the purpose of studying the relation of temperature and relative humidity to the production and dissemination of conidia, detailed life histories of a large number of individual spots on 10 plants in the medium-early field at Rocky Ford were kept during the season of 1913. The temperature and humidity records used in these correlations were those taken among the beet leaves near the ground and, together with rainfall and dates of irrigation, are shown in figure 7.

Beginning with the outermost or oldest, the leaves were tagged and numbered consecutively, and the location of the spots on each was indi-

cated on diagrams. As new leaves developed, they were included in the observations, and this was true also of new spots, until they became too numerous, after which only a few representative ones on each leaf were studied in detail. During the period from the 24th of June to the 19th of September 330 spots were studied, both surfaces being examined at frequent intervals with a hand lens. For the purpose of getting a basis for comparison of rates of development at different stages in the life history of the disease, percentage values were assigned to each stage as follows, the spots being grouped and averaged later (Table IV):

Percentage value.	Stage of development of fungus.
5.0	Spot first noticed. Neither conidia nor conidiophores present.
12.5	Conidiophores present.
19.7	Very few conidia.
25	Few conidia.
31.2	Conidia fairly numerous.
37.5	Conidia numerous.
43.7	Conidia fairly abundant.
50	Conidia abundant.

The value of the spot is the sum of the values of the two sides—that is the value of a spot on which there were but few conidia (25) on one side, and abundant conidia (50) on the other, is 75. Again, the value of a spot on which conidiophores only (12.5) were present on one side, and very few conidia (19.7) on the other, is 32.<sup>a</sup>

TABLE IV.—Data on life histories of a representative number of leaf spots studied on 10 plants in a medium-early sugar-beet field near Rocky Ford, Colo., during the season of 1913<sup>b</sup>

Plant No.	Leaf No.	Spot No.	Date.	Graph values.	Data on life histories.
*2	3	5	July 7, 1913.	10	No conidiophores on either surface. July 8, no change.
				37	Conidiophores above, conidia few below.
				50	Conidia few on both surfaces. July 14, no change.
				50	Conidia very few above, fairly numerous below.
				100	Conidia abundant on both surfaces. July 23, no change.
				62	Conidia fairly numerous on both surfaces.
				37	Conidia none above and few below.
2	4	1	July 7, 1913.	10	No conidiophores on either surface. July 7, no change.
				8	Conidiophores on both surfaces.
				69	Conidia very few above and abundant below.
				12	Conidia few above and abundant below.
				100	Conidia abundant on both surfaces. July 21, 23 (leaf yellow), July 25, no change.
				63	Conidia numerous above and few below.
				100	Conidia abundant on both surfaces.
2	4	3	July 7, 1913.	10	No conidiophores on either surface. July 8, no change.
				25	Conidiophores on both surfaces.
				62	Conidia fairly numerous on both surfaces.
				82	Conidia numerous above and fairly abundant below.
				100	Conidia abundant on both surfaces.
				100	Conidia abundant on both surfaces.
				100	Conidia abundant on both surfaces.
2	4	4	July 7, 1913.	10	No conidiophores on either surface.
				25	Conidiophores on both surfaces.
				10	Conidia few above and very abundant below. July 14, no change.
				57	Conidia very few above, numerous below.
				57	Conidia very few above, numerous below.
				57	Conidia very few above, numerous below.
				57	Conidia very few above, numerous below.

<sup>a</sup> For convenience the decimal fractions, which make only a negligible difference in the averages, are omitted.

<sup>b</sup> In Table IV asterisks (\*), daggers (†), and section marks (§) are used to designate definite leaf spots to which reference is made in the text.

TABLE IV.—Data on life histories of a representative number of leaf spots studied on 10 plants in a medium-early sugar-beet field near Rocky Ford, Colo., during the season of 1913—Continued

Plant No.	Leaf No.	Spot No.	Date.	Graph values.	Data on life histories.
** <sub>2</sub>	6	1	1913. July 7	15	Conidiophores forming above, nothing below.
			8	31	Conidiophores above, conidia very few below.
			10	100	Heavy production of conidia on both surfaces. July 12, 14, 16, 21, 23, no change.
			25	63	Conidia few above and numerous below.
3	2	1	June 24	10	No conidiophores on either surface. June 25, 26, 27, 28, 30, July 1, 2, 7 (leaf yellow), 8, no change.
3	2	2	25	10	No conidiophores on either surface. June 25, 26, 27, 28, 30, July 1, 2, 7, 8, no change.
3	4	1	July 7	10	No conidiophores on either surface.
				15	Conidiophores forming only on lower surface.
				62	Conidiophores abundant above and conidia abundant below.
				100	Conidia abundant on both surfaces. July 14, no change.
				16	Conidia very few above and abundant below.
				21	No conidia on either surface.
** <sub>3</sub>	8	1		7	No conidiophores on either surface. July 8, no change.
				10	Conidiophores abundant on both surfaces.
				12	Conidia abundant on both surfaces. July 14, 16, 21 (center of spot gone), no change.
				23	Conidia fairly abundant above and abundant below.
				25	Conidia few on both surfaces, leaving and leaf yellowing.
** <sub>3</sub>	8	2		7	No conidiophores on either surface. July 8, no change.
				10	Conidiophores abundant on both surfaces.
				12	Conidia abundant on both surfaces. July 14, no change.
				16	Conidia fairly abundant above and abundant below.
				21	Conidia abundant on both surfaces. July 23, no change.
				25	Conidia few on both surfaces.
† <sub>3</sub>	8	3		7	No conidiophores on either surface. July 8, no change.
				10	Conidiophores abundant on both surfaces.
				12	Conidia abundant on both surfaces. July 14, no change.
				16	Conidia fairly abundant above and abundant below.
				21	Conidia abundant on both surfaces. July 23, no change.
3	9	1		25	Conidia few above and abundant below.
				26	Conidiophores numerous above and few below.
				28	Conidia few above and very few below.
				50	Conidia few and matted together on both surfaces.
4	1	1	June 24	10	No conidiophores on either surface. June 25 (leaf dying), 26 (leaf dead), 27, 28, 30, July 1, no change.
4	3	1	July 8	10	No conidiophores on either surface.
				12	Conidia fairly numerous on both surfaces.
				14	Conidia leaving, few on both surfaces. July 16, 21 (leaf dead), no change.
** <sub>4</sub>	5	1	June 30 July 7	10	No conidiophores on either surface. July 1, 2, no change.
				30	Conidia forming on both surfaces. July 8, no change.
				25	Only conidiophores on both surfaces.
				100	Conidia abundant on both surfaces. July 14, no change.
				88	Conidia fairly abundant on both surfaces. July 21, 23, no change.
				25	Conidia few on both surfaces.
* <sub>4</sub>	12	1		50	Conidia few on both surfaces. July 23, no change.
				76	Conidia numerous on both surfaces.
5	1	2	1	10	No conidiophores on either surface. July 2, 7, 8, no change.
5	2	1		2	No conidiophores on either surface. July 7, no change.
				8	Conidiophores forming only on upper surface.
				10	Conidiophores above and conidia forming below.
				12	Conidia few on both surfaces.
				14	Conidia very few above, and fairly numerous below.
				16	Leaf dead, no change in spot.

TABLE IV.—Data on life histories of a representative number of leaf spots studied on 10 plants in a medium-early sugar-beet field near Rocky Ford, Colo., during the season of 1913—Continued

Plant No.	Leaf No.	Spot No.	Date.	Graph values.	Data on life histories.
†5	4	3	1913- July 7	20	No conidiophores above, very few conidia forming below.
			8	37	Conidiophores on upper surface, few conidia on lower.
			10	88	Conidia numerous above and very abundant below.
			19	75	Conidia few above and abundant below. July 14, no change.
			21	100	Conidia abundant on both surfaces. July 23 (leaf yellow), July 25, no change.
5	4	5	28	63	Conidia few above and numerous below.
			8	25	Conidiophores on both surfaces.
			10	37	Conidia few above and conidiophores below. July 12, no change.
			14	50	Conidia few on both surfaces.
			21	100	Conidia abundant on both surfaces. July 23, no change.
**5	6	1	25	50	Conidia few on both surfaces.
			28	25	No conidia on either surface.
			7	17	Conidiophores none above and forming below.
			8	25	Conidiophores on both surfaces.
			10	75	Conidia few above and very abundant below. July 12, 14, 16, no change.
5	8	4	21	100	Conidia abundant on both surfaces. July 23, no change.
			25	57	Conidia very few above and numerous below.
			28	37	Conidia none above and few below.
			14	10	No conidiophores on either surface.
			16	17	Conidiophores none above and forming below.
*5	8	5	21	100	Conidia abundant on both surfaces. July 23, no change.
			25	75	Conidia abundant above and few below.
			23	25	Conidiophores on both surfaces.
			25	75	Conidia numerous on both surfaces.
5	10	1	7	40	Conidia very few on either surface but more on lower. July 8, no change.
			10	75	Conidia few above and abundant below.
			12	100	Conidia abundant on both surfaces. July 14, 16, 21, no change.
			7	35	Conidiophores forming above and conidia few below. July 8, no change.
			20	62	Conidiophores well developed above and conidia abundant below.
5	10	3	12	70	Conidia very few above and abundant below.
			14	75	Conidia few above and abundant below. July 16, no change.
			21	100	Conidia abundant on both surfaces. July 23, no change.
			14	10	No conidiophores on either surface.
			16	17	Conidiophores above and none below.
*5	12	1	21	100	Conidia abundant on both surfaces. July 23, no change.
			25	63	Conidia few above and numerous below.
			23	25	Conidiophores on both surfaces.
			25	70	Conidia very few above and abundant below.
			28	63	Conidia few above and numerous and matted below.
6	1	1	June 24	10	No conidiophores on either surface.
			25	70	Conidiophores very few above and abundant below. June 26, no change.
			27	31	Conidiophores above and conidia very few below. June 28, 30, July 1, 2 (leaf dying), no change.
			July 7	25	Conidiophores, but no conidia present.
7	6	1	7	10	No conidiophores on either surface.
			10	50	Conidia few above and fairly numerous below.
			19	63	Conidia none above and abundant below.
			12	100	Conidia abundant on both surfaces. July 14, 16, 21, 23, 25, no change.
			10	15	Conidiophores forming on upper surface only.
7	6	3	19	75	Conidia abundant above and few below.
			14	50	Conidia few on either surface. July 16, no change.
			21	100	Conidia abundant on both surfaces. July 23, 25, no change.
			16	10	No conidiophores on either surface.
			21	100	Conidia abundant on both surfaces. July 23, 25, no change.

TABLE IV.—Data on life histories of a representative number of leaf spots studied on 10 plants in a medium-early sugar-beet field near Rocky Ford, Colo., during the season of 1913—Continued

Plant No.	Leaf No.	Spot No.	Date.	Graph values.	Data on life histories.
*7	6	8	1913, July 21	55	Conidia few above and fairly numerous below. July 23, no change.
			25	100	Conidia abundant on both surfaces.
*7	8	5	21	37	Conidia very few on both surfaces. July 23, no change.
			25	75	Conidia numerous on both surfaces.
			28	50	Conidia few on both surfaces. July 30, no change.
7	8	5	21	50	Conidia few on both surfaces.
			23	26	Conidia few above and fairly numerous below.
			25	63	Conidia numerous above and few below.
			28	50	Conidia few on both surfaces. July 30, Aug. 1, no change.
*7	8	7	July 23	15	Conidiophores none above and few below.
			25	75	Conidia few above and abundant below. July 28, 30, no change.
			Aug. 1	50	Conidia none above and numerous below.
7	8	12	July 25	25	Conidiophores on both surfaces.
			28	45	Conidia few forming on both surfaces.
			30	50	Conidia few on both surfaces. August 1, no change.
7	15	3	28	25	Conidiophores on both surfaces.
			30	63	Conidia few above and numerous below.
			Aug. 1	88	Conidia numerous above and abundant below.
†††8	8	5	July 9	70	No conidiophores on either surface.
			10	100	Conidia abundant on both surfaces.
*8	12	1	23	50	Conidia few on both surfaces.
			25	75	Conidia numerous on both surfaces. July 28, no change.
			30	63	Conidia numerous above and abundant below. August 1, no change.
			Aug. 5	75	Conidia numerous on both surfaces. Aug. 7, 9, 11, no change.
8	23	13	5	15	Conidiophores few above and none below. Aug. 7, no change.
			9	25	Conidiophores on both surfaces.
			11	57	Conidia numerous above and very few below.
			13	100	Conidia abundant on both surfaces.
†8	23	18	7	10	No conidiophores on either surface. Aug. 9, no change.
			11	60	Conidia abundant above and conidiophores few below.
			13	63	Conidia numerous above and few below. Aug. 15, 18, no change.
			22	100	Conidia abundant on both surfaces.
†8	24	1	July 28	75	Conidia numerous on both surfaces.
			30	63	Conidia fairly numerous on both surfaces. Aug. 1, no change.
			Aug. 5	44	Conidia few above and very few below.
8	25	3	7	75	Conidia numerous and matted above and numerous below.
			1	10	No conidiophores on either surface.
			5	44	Conidia very few above and few below.
*8	29	1	7	56	Conidia few above and fairly numerous below.
			1	10	No conidiophores on either surface.
			5	63	Conidia fairly numerous on both surfaces.
			7	75	Conidia numerous on both surfaces.
			9	50	Conidia few on both surfaces.
			11	50	Conidia few above and numerous below.
			13	100	Conidia abundant on both surfaces. Aug. 15, no change.
			18	75	Conidia few above and abundant below.
			22	88	Conidia numerous above and abundant below. Aug. 25, 28, no change.
			30	63	Conidia few above and numerous below. Sept. 1, no change except conidia matted above.
8	29	3	Sept. 3	57	Conidia very few above and numerous below.
			Aug. 5	25	Conidiophores on both surfaces.
			7	30	Conidiophores above and conidia forming below.
			9	50	Conidia very few above and fairly numerous below.
			11	100	Conidia abundant on both surfaces. Aug. 13, 15, 18, 22, 25, no change.

TABLE IV.—Data on life histories of a representative number of leaf spots studied on 10 plants in a medium-early sugar-beet field near Rocky Ford, Colo., during the season of 1913—Continued

Plant No.	Leaf No.	Spot No.	Date.	Graph values.	Data on life histories.
***†8	29	6	1913- Aug. 5	20	Conidiophores very few on both surfaces. Aug. 7, no change.
			Aug. 9	75	Conidia numerous on both surfaces. Aug. 11, no change.
			Aug. 13	100	Conidia abundant on both surfaces. Aug. 15, no change.
			Aug. 18	75	Conidia few above and abundant below.
			Aug. 22	69	Conidia fairly numerous above and numerous below.
			Aug. 25	100	Conidia abundant on both surfaces.
			Aug. 28	87	Conidia fairly abundant on both surfaces.
			Sept. 30	75	Conidia numerous on both surfaces.
			Sept. 1	69	Conidia fairly numerous above and numerous below.
			Sept. 3	62	Conidia fairly numerous and matted above and fairly numerous below.
***8	29	8	Aug. 5	10	No conidiophores on either surface. Aug. 7, 9, no change.
			Aug. 11	24	Conidia very few above and nothing below.
			Aug. 13	88	Conidia abundant above and numerous below.
			Aug. 15	64	Conidia fairly abundant above and abundant below.
			Aug. 18	88	Conidia numerous above and abundant below.
			Aug. 22	100	Conidia abundant on both surfaces.
			Aug. 25	81	Conidia abundant above and fairly numerous and matted below.
8	29	9	Aug. 5	35	Conidiophores few above and conidia few below.
			Aug. 7	70	Conidiophores numerous above and conidia numerous below.
			Aug. 9	88	Conidia numerous above and abundant below.
			Aug. 11	37	Conidia few above and none below.
			Aug. 13	75	Conidia numerous and matted above and numerous below.
			Aug. 15	75	Aug. 15, no change.
			Aug. 18	88	Conidia numerous above and abundant below.
***†8	29	11	Aug. 7	10	Conidiophores few on both surfaces.
			Aug. 9	44	Conidia very few above and few below.
			Aug. 13	75	Conidia numerous on both surfaces. Aug. 15, no change.
			Aug. 18	88	Conidia numerous above and abundant below.
			Aug. 22	63	Conidia very few and matted above and fairly abundant below.
			Aug. 25	88	Conidia numerous and matted above and abundant below.
			Aug. 28	75	Conidia few above and abundant below.
			Sept. 30	88	Conidia numerous and matted above and abundant below.
			Sept. 1	82	Conidia numerous and matted above and fairly abundant below.
§8	35	1	Aug. 5	69	Conidia few and matted above and fairly abundant below.
			Aug. 7	50	Conidia very few and matted above and fairly numerous and matted below.
			Aug. 8	43	Conidia none above and fairly numerous and matted below.
††8	35	9	Aug. 1	10	No conidiophores on either surface.
			Aug. 5	81	Conidia fairly numerous and matted above and abundant below. Aug. 7, no change.
			Aug. 9	69	Conidia fairly numerous and matted above and numerous below.
			Aug. 11	63	Conidia numerous above and few below.
			Aug. 13	100	Conidia abundant on both surfaces.
			Aug. 15	81	Conidia fairly numerous and matted above and abundant below.
			Aug. 18	63	Conidia few and matted above and numerous below. Aug. 25, no change.
			Aug. 28	50	Conidia none above and numerous below.
			Sept. 30	36	Conidia none above and fairly abundant below.
††8	35	9	Aug. 11	10	No conidiophores on either surface.
			Aug. 15	15	Conidiophores few above and none below.
			Aug. 18	37	Conidia very few on both surfaces.
			Aug. 22	44	Conidia very few above and few below.
			Aug. 25	62	Conidia fairly numerous on both surfaces.
			Aug. 28	69	Conidia numerous above and fairly numerous below. Aug. 30, Sept. 1, no change.
			Sept. 3	62	Conidia fairly numerous and matted above and fairly numerous below.
			Sept. 6	30	Conidia few and matted on both surfaces.
			Sept. 8	37	Conidia very few on both surfaces. Sept. 11, no change except matted on both surfaces.

TABLE IV.—Data on life histories of a representative number of leaf spots studied on 10 plants in a medium-early sugar-beet field near Rocky Ford, Colo., during the season of 1913—Continued

Plant No.	Leaf No.	Spot No.	Date.	Graph values.	Data on life histories.
H. 88	35	10	1913.		
			Aug. 13	10	No conidiophores on either surface.
			15	15	Conidiophores very few above and none below. August 18, no change.
			22	37	Conidia very few on both surfaces.
			25	50	Conidia fairly numerous above and few below.
			28	50	Conidia fairly numerous above and very few below.
			30	88	Conidia numerous and matted above and abundant below.
			Sept. 1, 5		no change.
			8	50	Conidia few and matted on both surfaces.
			10	37	Conidia very few and matted on both surfaces.
H. 88	37	1	Aug. 11	24	Conidia very few above and nothing below.
			13	100	Conidia abundant on both surfaces. August 15, no change except conidia matted above.
			18	94	Conidia fairly abundant and matted above and abundant below.
			22	88	Conidia numerous above and abundant below. August 25, no change.
			28	75	Conidia numerous on both surfaces.
			30	69	Conidia fairly numerous above and numerous below. September 1, 3, no change.
			Sept. 6	37	Conidia very few on both surfaces.
			8	75	Conidia none on either surface. September 10, 13, 15, no change.
H. 88	37	2	Aug. 11	10	No conidiophores on either surface.
			13	100	Conidia abundant above and abundant and matted below.
			15	94	Conidia abundant above and fairly abundant below.
			18	87	Conidia fairly abundant on both surfaces.
			22	100	Conidia abundant on both surfaces. August 25, no change except slightly matted on both surfaces.
			28	94	Conidia abundant and matted above and fairly abundant below.
			30	100	Conidia abundant and matted on both surfaces.
			Sept. 1	87	Conidia fairly abundant and matted on both surfaces. September 3, 5, no change.
			8	57	Conidia very few above and fairly numerous below. September 10, no change.
			13	37	Conidia none above and few below. September 15, no change. Conidia still matted on both surfaces.
H. 88	37	3	Aug. 11	10	No conidiophores on either surface.
			13	18	Conidiophores very few on both surfaces.
			15	44	Conidia very few above and few below.
			18	50	Conidia very few above and fairly numerous below.
			22	62	Conidia fairly numerous and matted above and fairly numerous below.
			25	62	Conidia fairly numerous on both surfaces.
			28	75	Conidia numerous on both surfaces. August 30, no change. Conidia matted on both surfaces.
			Sept. 1	62	Conidia fairly numerous and matted on both surfaces.
88	37	6	Aug. 18	10	No conidiophores on either surface.
			22	50	Conidia few on both surfaces.
			25	62	Conidia fairly numerous on both surfaces.
			28	75	Conidia fairly abundant in center on both surfaces.
			30	70	Conidia numerous in center above and fairly abundant in center below. September 1, no change.
			Sept. 3	69	Conidia fairly numerous above and fairly abundant in center below.
			8	87	Conidia fairly abundant on both surfaces.
			13	75	Conidia fairly numerous and matted above and fairly abundant below. September 10, no change.
			17	94	Conidia fairly abundant above and abundant below. September 15, no change.
				50	Conidia few on either surface.
8	38	1	Aug. 11	10	No conidiophores on either surface.
			13	69	Conidia numerous above and fairly numerous below.
			15	69	Conidia fairly numerous above and numerous and matted below.
			18	82	Conidia fairly abundant above and numerous and matted below.
			22	69	Conidia numerous and matted above and fairly numerous below.
8	39	6	Aug. 18	10	No conidiophores on either surface.
			22	63	Conidia numerous above and few below.
			25	82	Conidia fairly abundant above and numerous below.

TABLE IV.—Data on life histories of a representative number of leaf spots studied on 10 plants in a medium-early sugar-beet field near Rocky Ford, Colo., during the season of 1913—Continued

Plant No.	Leaf No.	Spot No.	Date.	Graph values.	Data on life histories.
8	41	3	1913.		
			Aug. 22	31	Conidiophores above and conidia very few below.
			25	62	Conidia fairly numerous on both surfaces.
			28	82	Conidia numerous above and fairly abundant below.
			30	88	Conidia numerous above and abundant below. September 7, no change.
			Sept. 3	74	Conidia fairly numerous above and fairly abundant below.
			6	69	Conidia fairly numerous and matted above and numerous and matted below. September 8, 10, no change.
			13	57	Conidia very few and matted above and numerous below.
			19	50	September 13, 17, no change. Conidia none above and numerous below.
††, §8	44	1	Aug. 25	10	No conidiophores on either surface.
			28	74	Conidia very few above and nothing below.
			30	44	Conidia few above and very few below. September 1, 3, 6, no change.
			Sept. 8	50	Conidia few on both surfaces.
			10	50	Conidia fairly numerous above and few below.
			13	41	Conidia very few above and few below.
			15	69	Conidia fairly numerous and matted above and numerous below.
			17	44	Conidia very few and matted above and few below.
			19	37	Conidia none above and very few below.
††8	44	2	Aug. 25	15	Conidiophores forming above and nothing below.
			28	75	Conidia fairly abundant in center of both surfaces.
			30	90	Conidia abundant in center of both surfaces. September 1, 3, 6, no change.
			Sept. 8	69	Conidia fairly numerous above and numerous below.
			10	100	Conidia abundant on both surfaces. September 13, 15, no change.
			17	56	Conidia few and matted above and fairly numerous and matted below.
			19	50	Conidia very few and matted above and fairly numerous and matted below.
8	44	3	Aug. 25	10	No conidiophores on either surface.
			28	80	Conidia fairly abundant in center on both surfaces.
			30	90	Conidia fairly abundant in center above and abundant in center below. September 1, 3, no change.
			Sept. 6	87	Conidia fairly abundant and matted above and fairly abundant below. September 8, no change.
			10	94	Conidia fairly abundant above and abundant below. September 13, no change.
			13	88	Conidia numerous and matted above and abundant below.
			17	75	Conidia fairly numerous and matted above and fairly abundant below.
			19	61	Conidia few and matted above and numerous below.
††, §8	45	1	Aug. 22	37	Conidia very few on both surfaces.
			25	39	Conidia few above and fairly numerous and matted below.
			28	82	Conidia numerous and matted above and fairly abundant below.
			30	88	Conidia numerous above and abundant below. September 1, no change.
			Sept. 3	81	Conidia fairly numerous and matted above and abundant below. September 6, 8, no change except conidia matted below.
			10	75	Conidia fairly numerous and matted above and fairly abundant and matted below.
			13	69	Conidia few and matted above and fairly abundant and matted below.
			15	50	Conidia none above and numerous and matted below.
			17	31	Conidia none above and very few and matted below.
			19	25	Conidia none on either surface.
8	45	2	Aug. 21	10	No conidiophores on either surface.
			28	45	Conidia few above and conidiophores forming below.
			30	25	Conidia very few above and nothing below.
			Sept. 1	29	Conidia very few above and few conidiophores below. September 3, no change.
			6	44	Conidia few above and very few below.
			8	50	Conidia few and matted above and few below.
			10	44	Conidia few and matted above and very few below. September 13, 15, 17, no change.
			19	25	Conidia none on either surface.



Plant No.	Leaf No.	Spot No.	Date.	Graph values.	Data on life histories.	
8	45	5	1912. Aug. 25	10	No conidiophores on either surface.	
			28	56	Conidia few above and fairly numerous below. August 30, September 7, 3, no change.	
			Sept. 6	65	Conidia few and matted above and fairly abundant in center below.	
			8	58	Conidia few and matted above and numerous in center below.	
			10	69	Conidia few and matted above and fairly abundant below.	
			13	63	Conidia few and matted above and numerous in center below. September 15, no change.	
8	49	1	Aug. 28	37	Conidia very few on both surfaces.	
			30	50	Conidia few on both surfaces. September 1, 3, no change.	
			Sept. 6	50	Conidia numerous above and fairly numerous below.	
			8	62	Conidia fairly numerous and matted above and fairly numerous below.	
			10	75	Conidia numerous and slightly matted on both surfaces.	
			13	56	Conidia fairly numerous and matted above and few below.	
8	49	2	Aug. 28	10	No conidiophores on either surface.	
			30	25	No conidiophores above and conidia very few below. September 3, no change.	
			Sept. 6	56	Conidia few above and fairly numerous below.	
			8	50	Conidia very few above and fairly numerous below. September 10, 13, no change.	
8	49	3	Aug. 28	10	No conidiophores on either surface.	
			30	15	Conidiophores few above and very few below.	
			Sept. 1	30	Conidiophores few above and conidia very few below. September 3, no change.	
			6	62	Conidia fairly numerous on both surfaces.	
			8	65	Conidia fairly numerous above and numerous in center below.	
			10	75	Conidia numerous above and numerous and matted in center below.	
			13	62	Conidia fairly numerous on both surfaces.	
8	49	6		1	No conidiophores upon either surface. September 3, no change.	
				6	44	Conidia very few above and few below. September 8, no change.
				10	50	Conidia few on both surfaces. September 13, no change.
8	49	7		1	10	No conidiophores on either surface. September 3, no change.
				6	24	Nothing above and conidia very few below.
				8	44	Conidia very few above and few below. September 10, 13, no change.
8	49	8		1	10	No conidiophores on either surface. September 3, 6, no change.
				8	20	Nothing above and conidia forming below.
				10	44	Conidia very few above and few below. September 13, no change.
8	49	9		3	10	No conidiophores on either surface. September 6, no change.
				8	20	Nothing above and very few conidia forming below.
				10	44	Conidia very few above and few below. September 13, no change.
8	49	10		3	10	No conidiophores on either surface. September 6, 8, no change.
				10	37	Conidia very few on both surfaces. September 13, no change.
8	51	1		3	10	No conidiophores on either surface. September 6, no change.
				8	24	Conidia very few above and nothing below. September 10, no change.
				13	37	Conidia very few upon both surfaces. September 15, 17, no change.
8	51	2		6	15	No conidiophores on either surface.
				8	27	Conidia very few on both surfaces.
				10	65	Conidia numerous in the center upon both surfaces. September 13, 15, no change.
				17	44	Conidia very few above and few below.
8	51	3		6	10	No conidiophores on either surface. September 8, no change.
				10	37	Conidia very few on both surfaces.
				13	44	Conidia few above and very few below. September 15, 17, no change.

TABLE IV.—Data on life histories of a representative number of leaf spots studied on 10 plants in a medium-early sugar-beet field near Rocky Ford, Colo., during the season of 1913—Continued

Plant No.	Leaf No.	Spot No.	Date.	Graph values.	Data on life histories.
8	51	4	1913- Sept. 6	10	No conidiophores on either surface. September 8, 10, no change.
			13	30	Conidiophores forming above and conidia very few below. September 15, 17, no change.
8	51	5	6	10	No conidiophores on either surface. September 8, 10, 13, 15, no change.
				15	No conidiophores on either surface. September 8, 10, 13, no change.
				24	Conidiophores none above and forming below. Nothing above and very few conidia below.
8	53	1	17	10	No conidiophores on either surface. September 19, no change.
				17	No conidiophores on either surface. September 19, no change.
9	3	1	June 24	10	No conidiophores on either surface. June 25, 26, 27, 28, 30; July 1, 2, 7, no change.
				10	No conidiophores on either surface.
9	3	2	July 9	10	Conidiophores abundant on both surfaces.
				25	Conidia fairly abundant on both surfaces. July 14, no change.
				100	Conidia abundant on both surfaces.
				37	Leaf dead, conidia very few on both surfaces. July 25, 28, no change.
9	5	1	21	10	Conidia few on both surfaces. July 23, 25, no change.
				37	Conidia none above and few below. July 30, no change.
9	5	2	21	17	Conidiophores above and none below.
				75	Conidia numerous on both surfaces. July 28, 30, no change.
9	7	1	9	10	No conidiophores on either surface.
				37	Conidia few above and conidiophores abundant below.
				94	Conidia abundant above and fairly abundant below. July 14, 26, no change.
10	3	1	June 30	10	No conidiophores on either surface. July 1, 2, 7, 9, 10, 11, no change.
				10	No conidiophores on either surface.
10	5	1	July 2	10	Conidiophores few above and abundant below.
				29	Conidiophores above and conidia few below.
				75	Conidia numerous on both surfaces. July 15, 24, no change.
				100	Conidia abundant on both surfaces. July 21, 23, no change.
				75	Conidia few above and abundant below.
				50	Conidia none above and numerous below. Leaf dead. July 30, no change.
10	5	3	21	50	Conidia few on both surfaces. July 23, no change.
				50	Conidia fairly numerous above and few below.
				25	Conidia none on either surface. July 30, no change.
10	5	4	21	37	Conidiophores above and conidia few below.
				44	Conidia very few above and few below.
				37	Conidia very few on both surfaces.
				25	Conidia none on either surface. July 30, no change.
10	6	1	2	10	No conidiophores on either surface.
				37	Conidia very few above and conidiophores below.
				62	Conidia fairly numerous on both surfaces. July 10, 12, 14, no change.
				50	Conidia few above and fairly numerous below.
10	6	6	July 12	10	No conidiophores on either surface.
				15	Conidiophores none above and very few below.
				37	Conidiophores above and conidia few below.
				100	Conidia abundant on both surfaces. July 23, 25, 28, 30, no change.

TABLE IV.—Data on life histories of a representative number of leaf spots studied on 10 plants in a medium-early sugar-beet field near Rocky Ford, Colo., during the season of 1913.—Continued

Plant No.	Leaf No.	Spot No.	Date.	Graph values.	Data on life histories.
**10	9	1	1913, July 7	30	Conidia few on both surfaces. July 9, no change.
			10	75	Conidia few above and abundant below.
			12	94	Conidia fairly abundant above and abundant below. July 14, no change.
			16	69	Conidia very few above and abundant below.
			21	100	Conidia abundant on both surfaces. July 23, no change.
			25	75	Conidia numerous on both surfaces.
			7	37	Conidiophores above and conidia few below. July 9, no change.
			10	75	Conidia few above and abundant below.
			12	69	Conidia few above and fairly abundant below. July 14, no change.
†10	9	2	16	69	Conidia very few above and abundant below.
			21	100	Conidia abundant on both surfaces. July 23, 25, no change.
10	9	3	23	25	Conidiophores on both surfaces.
			25	75	Conidia numerous on both surfaces. July 28, no change.
*10	10	1	23	10	No conidiophores on either surface.
			25	75	Conidia numerous on both surfaces.
			28	75	Conidia abundant above and few and matted below. July 30, no change.
10	11	1	25	75	Conidia numerous on both surfaces.
			28	75	Conidia abundant above and few and matted below. July 30, no change.
10	11	3	25	50	Conidiophores above and conidia numerous below.
			28	75	Conidia numerous on both surfaces. July 30, no change.




FIG. 6.—Curves of the leaf spot history series, showing the production of conidia on different dates from June 24 to September 19, 1913, at Rocky Ford, Colo.

After the values were all assigned, the spots which appeared on or about the same day were brought together in 24 groups and averaged (Table V and fig. 6). The temperature and humidity records used in the correlations with the leafspot histories were taken among the sugar-beet leaves near the surface of the ground.

TABLE V.—Leafspot history series showing their arbitrary values on different dates and the number of spots entering into the average of each series at Rocky Ford, Colo., 1913

[illegible]

**CONIDIAL PRODUCTION.**—Under favorable conditions conidia are produced apparently much more readily by young than by old leaf-spots. For instance, as will be seen in Table IV, leaf spots (\*)<sup>1</sup> 2 to 4 days old showed a marked increase in conidial production from July 23 to 25, while during the same period spots (\*\*) 14 to 24 days old in most cases showed a decrease. However, spots (\*\*\*) 2 to 3 weeks old on green leaves showed an increased production from August 18 to 25, the conditions being favorable, and some (†) even produced a second and third crop, although usually but one crop (††) is produced and this while the spots are comparatively young. It was also found that under favorable conditions a spot (†††) may produce abundant conidia on both surfaces in one day. Usually the maximum production is reached within 10 days after the spots appear (fig. 6), and sometimes under very favorable conditions the production may increase after this period (fig. 6, curves D and E, July 17 to 23), but the older spots do not always respond to favorable conditions in this way (fig. 6, curves C and F). In no case was a new growth of conidia observed on spots on yellow or dying leaves on green plants in the field. The fungus seemed to lose its vigor much sooner on such leaves than on green leaves which remained attached to the crown at harvest time. From the standpoint of control of the disease this is a very important point, from the fact that at harvest time the green leaves, on which the fungus is vigorous, are removed with the crowns and stored in the silo, while the yellow and dying leaves, on which the fungus may be too weak to overwinter, break off and remain on the ground.

During the greater part of August and September, when the precipitation was light (fig. 7), many of the conidia had a shrunken appearance and were massed together on the leafspot areas (Table IV, §). When placed in water, these conidia did not germinate; consequently this desiccation of the conidia may also be an important factor in connection with the vitality of the fungus on the host.

The position of the leaf on which the spot studied was located was also found to be an important factor in conidial production, an abundance of conidia being frequently observed on leaves protected from the sun, while at the same time few were observed on those exposed to the sun the greater part of the day. This difference in production is thought to be due mainly to the difference in humidity of the protected and the exposed locations.

A study of the comparative production of conidia on the upper and the lower surfaces of the spots was also made, the conidia on the spots included in series E, K, N, and S (fig. 6) being tabulated for this purpose.

Generally a more abundant conidial production was found on the lower than on the upper surface (fig. 8), and this was due apparently to

<sup>1</sup> The asterisks (\*), daggers (†), and section marks (§) refer to particular leaf spots in Table IV.

the probably higher humidity of the former. Only during a very favorable period (fig. 7, July 19 to 21) or where the leaves were turned up or protected by other leaves was the conidial production on the upper surface equal to that on the lower surface (fig. 8, series E, July 21). At times, conidia were formed more abundantly on the upper surface than on the lower (fig. 6, series N, August 11, and series S, August 28). Because of the spongy parenchyma and the greater number of stomata on the lower surface, it might be supposed that conidiophores could be produced more readily on this than on the upper surface; but, as above indicated, humidity would seem to be the controlling factor in this connection.




FIG. 7.—Curves of the maximum and minimum temperatures and humidities, the number of hours that the humidity remained above 60 from noon of the preceding to noon of the given day among the plants, and rainfall and irrigation records, taken in a medium-early sugar-beet field from June 20 to September 22, 1913, at Rocky Ford, Colo.

A comparison of the conidial production as shown in Table IV and figure 6 and the climatic data shown in figure 7 indicates many definite relations. When the spots were first found, on June 20 and 24, conidia were fairly numerous (fig. 6, curve A) on all except six spots, which had evidently just developed on the latter date, as no conidia were present at this time and conidiophores only were produced the next two weeks. The following week there was but little increase, and during the next few days many of the conidia were disseminated. The small production of conidia was evidently due directly to the high temperature and the low humidity which prevailed during this period (fig. 7), as conidia were produced in great abundance from July 9 to 12 (fig. 6, curves C, D, E, F), when the temperature was lower and the humidity higher (fig. 7). Dur-

ing this time and the few days just preceding, the humidity remained above 60 for a longer time on an average and the minimum humidity did not become so excessively low nor the temperature so excessively high as during the time previous to July 4.

The next period of pronounced increase in conidial production was from July 19 to 23 (fig. 6, curves D, E, C), when the conditions were more favorable than during any period of similar length through the summer, the humidity ranging above 60 on an average of 19.4 hours each day and not falling below 52 (fig. 7), and the temperature ranging from 60° to 90° F.

Conidial production was again above the average (curves M, N, and O) from August 9 to 13, during which period the humidity remained above 60 from 13 to 20 hours each day and there was a small amount of rain which seemed to aid in maintaining the necessary humid conditions. Production was checked on August 16, on which date the temperature was 102° and the average humidity low, and was again inhibited after




FIG. 8.—Curves of the comparative production of conidia on the upper and lower surfaces of the leaf spots, representing series E, K, N, and G of Table V and figure 6. Rocky Ford, Colo., 1913.

September 11, subsequent to which date the minimum temperatures ranged from about 30° to 45° and the maximum from about 65° to 83°, while the humidity remained above 60 for 12.4 hours per day, on an average.

The general conclusion from these tests is that conidial production is greatly influenced by temperature and relative humidity, or speaking specifically—

- (1) A temperature of 100° F. or over is detrimental to conidial production, directly perhaps because it is inimical to the growth of the fungus and indirectly because humidity is ordinarily excessively low at such an extreme temperature.
- (2) Conidial production is greatly checked at daily temperatures ranging below 50° as a minimum and 80° as a maximum.
- (3) The most favorable temperature for conidial production is 80° to 90° in the daytime and not below 60° at night.
- (4) The temperature being favorable, the largest conidial production occurred at the higher humidities. A good production occurred when

the humidity remained above 60 for not less than 15 to 18 hours, but very few were produced when the humidity remained above 60 for less than 10 to 12 hours daily.

With a view to determining the approximate number of conidia produced on a sugar-beet plant under a favorable temperature and humidity, one representing a heavy infection in August was selected. After the infected leaves were measured a representative portion of conidia were carefully washed off into sterilized water and counted. The count, which was made by means of a dilution method, showed 250,000,000 conidia on the plant at that time.

CONIDIAL DISSEMINATION.—That a period of low humidity, with its accompanying factors, is favorable to the dissemination of conidia was frequently observed (fig. 6, curve R). For instance, it was found that the amount of conidia diminished on September 1, 6, 14, and 15, when the humidity remained above 60 for 5, 6, 10, and 4 hours, respectively; while, on the other hand, there was no diminution in the amount present on September 3 to 5 and 8 to 10, during which periods the humidity remained above 60 for 12 to 16 hours.

Rainfall is also an important factor in the dissemination of conidia, as was noted in several instances. On July 19 (fig. 6, curve F) rain fell, and as a result many conidia were washed off, and the same was true in the case of rains on July 23 (curves C, D, E, G, H), August 9 (curve K), September 4 (curves N, O, R), and September 16 (curves Q, S, T, U, V, W). After rains on July 19, August 9, and September 4, however, there were more conidia present than before, but this was probably due to the fact that more were produced under the favorable humid conditions attending these rains than were washed off. It was also found that the conidia were disseminated more rapidly from the upper than from the lower surface of the spots (fig. 8). This was due probably to the greater exposure of the former to wind and rainfall.

#### RELATION OF INFECTION CYCLES TO CLIMATIC CONDITIONS

For the purpose of determining the relation of infection cycles to climatic conditions, a study was made of the increase and spread of disease in a field of sugar beets planted about May 1<sup>1</sup> at Rocky Ford and one planted about two weeks later. Both fields had been in beets for two or three years, and as very few, if any, of the tops were removed after the harvest of 1912, infection appeared early in 1913 and was generally distributed.

Three plants in the early field (Table VII) and ten in the medium-early (Table VI) were selected, the leaves tagged and numbered consecutively, beginning with the outermost or oldest and continuing with the new ones as they appeared. The spots on each leaf were counted at

<sup>1</sup> Conidial production and dissemination were also studied in this field.



frequent intervals,<sup>1</sup> and the average actual increase of spots per plant computed (Table VIII and fig. 9). It was found that from 400 to 1,000 spots on a leaf, depending on its size, killed it within a few days.




FIG. 9.—Curves of the 2-day average increases in the number of leaf spots per plant in a medium-early and an early sugar-beet field, from June 18 to September 19, 1913, at Rocky Ford, Colo.

TABLE VI.—Average infection cycle of *Cercospora beticola* in a medium-early sugar-beet field with poorly developed foliage and with a consequent low humidity early in the season at Rocky Ford, Colo., in 1913

Date.	Total number of leaves marked.	Total number of leaves infected.	Total number of leaves dead.	Number of leaves killed by <i>Cercospora beticola</i> .	Number of infected green and dying leaves.	Number of functional leaves—		Total number of functional leaves.	Total number of leaf spots per plant.	Average number of leaf spots per leaf.
						Infected.	Uninfected.			
July 1.....	14.2	3.2	3.5		3	2.9	8	10.9	5	1.6
3.....	16.8	4.7	3.8		4.4	4.1	8.9	13	10.4	2.3
5.....	17.4	5.4	4.3		4.7	4.4	8.7	15.1	11.4	2.4
7.....	18.5	5.5	4.0		4.5	4.3	9.0	15.9	11.2	2.4
9.....	19.7	5.5	5.1		4.3	4	10.6	14.6	10.8	2.5
11.....	21.3	5.8	5.5		4.3	4	11.8	15.8	10.8	2.5
13.....	22.0	6	6.1		4.1	3.8	12.7	16.5	10.4	2.5
15.....	26.3	6.6	7.1		4.4	3.5	15.7	19.2	13.3	3
17.....	27.5	7.8	8		4.7	4.1	15.4	19.5	17.2	3.6
19.....	29	8.1	8.9		4.4	3.5	16.6	20.1	17.5	3.9
21.....	30.5	10.8	9.3		6.5	5.8	15.4	21.2	20.5	3.2
23.....	33	12.6	11		10.7	10.3	11.7	22	49	3.9
Aug. 2.....	34	27	12		20	17	3	22	93.5	4.7
4.....	35.5	27	12.5		19	18.5	4.5	23	292	12.7
6.....	36.5	27	13.5		18.5	17.5	5.5	24	355	16.1
8.....	37.5	29	13.5		19.5	19.5	4.5	24	402	20.6
10.....	38.5	30	15.5		20.5	19	4	21	455.5	22.2
12.....	40	31	15.5		20	20	4.5	24.5	593	29.6
14.....	41.5	31.5	15.5		20.5	20.5	5.5	26	875	41.9
16.....	43	33.5	15.5		22.5	22.5	5	27.5	1,155.5	54.3
18.....	45	35.5	15.5		24.5	24.5	5	29.5	1,704.5	85.9
20.....	46	38	15.5		27	27	3.5	30.5	3,120	170.6
22.....	48	41	17	1.5	29.5	25	3	31	9,681.5	307.8
24.....	48.5	41	18.5	2	28.5	25.5	3	30.5	10,756	380.9
Sept. 2.....	50	41.5	23.5	6	25.5	23.5	4	26.5	13,174.5	485.3
4.....	50.5	42.5	23.5	8	23.5	21.5	4	25	11,403	429
6.....	51.5	42.5	26.5	9	21.5	20.5	5	25	10,928.5	508.3
8.....	52.5	44.5	28.5	11	22.5	20.5	4.5	24	10,782	479.2
10.....	53	44.5	30	12.5	20.5	19	4	24	10,064.5	487.9
12.....	53.5	44.5	32.5	15	19.5	17.5	4.5	21	9,585.5	501.5
14.....	54	45.5	34.5	17	17.5	15.5	4	19.5	8,464	485.9
16.....	55	45.5	35	17.5	15.5	15	5	20	7,597.5	490.1

<sup>1</sup> For convenience and uniformity, 2-day averages were used in making the comparisons, the counts being made usually at 2-day intervals.

TABLE VII.—Average infection cycle of *Cercospora beticola* in an early sugar-beet field in which there was a heavy production of foliage and a consequent high humidity early in the season at Rocky Ford, Colo., in 1913

Date.	Total number of leaves marked	Total number of leaves infected.	Total number of leaves dead.	Number of leaves killed by <i>Cercospora beticola</i> .	Number of leaves infected green and dying leaves.	Number of functional leaves—		Total number of functional leaves.	Total number of leaf spots per plant.	Average number of leaf spots per leaf.
						Infected.	Uninfected.			
July 7	23	13.5	4	.....	12	12	7	19	539	44
9	24	13.5	4.5	.....	12	22.5	8	19.5	596.5	49.7
12	26	14	5.5	.....	12	11	9	20.5	615	51.2
14	27.5	14.5	5.5	.....	11.5	11.5	10.5	22	554.5	48.2
16	29.5	16	7	.....	11	11.5	10.5	22.5	574.5	44.2
21	34.5	23	8.5	.....	18.5	17	9	26	1,113.5	50.2
23	37	29	10.5	.....	23	21	5.5	26.5	2,216.5	96.3
25	38.5	30	11	.....	22	21.5	6	27.5	3,776.5	171.6
29	41.5	31	11.5	.....	22.5	22	8	30	7,045.5	373.1
Aug. 1	44	17	15	3	28	24.5	4.5	29	11,966.5	427.3
5	46.5	39	16	4	26.5	25.5	5	30.5	12,638	476.9
7	48.5	40	18	6	26.5	24.5	6	30.5	13,995	524.7
9	50	42	20	8	26.5	24.5	5.5	30	14,228.5	536.9
11	51.5	43	21.5	9.5	24.5	23.5	7	30	16,386	666.8
13	53.5	42.5	24.5	12.5	23.5	20.5	8.5	29	16,593	710.3
15	55.5	45	29.5	17.5	23	18	8	26	14,993	651.8

TABLE VIII.—Actual and 2-day average increase in the number of leaf spots per plant in a medium-early and an early sugar-beet field from June 18 to September 19, at Rocky Ford, Colo., in 1913

Date.	Increase in medium-early field.		Increase in early field.		Date.	Increase in medium-early field.		Increase in early field.	
	Actual.	2-day average.	Actual.	2-day average.		Actual.	2-day average.	Actual.	2-day average.
June 20	0.3	0.3	.....	.....	Aug. 5	203	102	3,996	1,998
22	0.5	0.5	.....	.....	7	65	65	2,107	2,107
24	0	0	.....	.....	9	127	127	2,373	2,373
26	0.3	0.3	.....	.....	11	53	53	4,208	4,208
28	1	1	.....	.....	13	190	190	1,831	1,831
30	0.4	0.4	.....	.....	15	282	282	1,450	1,450
July 2	2.3	2.3	.....	.....	18	280	280	2,459	1,513
7	4.9	1.9	.....	.....	22	295	447	2,286	1,743
8	1.4	2.8	.....	.....	25	1,379	919	2,282	1,722
9	.....	.....	70	70	28	5,064	3,776	2,944	1,664
10	0.9	0.9	.....	.....	30	2,572	2,572	.....	.....
12	0.2	0.2	18.5	12.2	1	2,582	2,582	.....	.....
14	0.7	0.7	24.5	24.5	3	1,614	1,614	.....	.....
16	0.7	0.7	20.5	20.5	6	799	533	.....	.....
21	3.3	1.3	622	209	8	771	771	.....	.....
23	3.3	3.3	1,222	1,372	10	578	578	.....	.....
25	1.9	1.9	1,723	1,723	13	712	488	.....	.....
28	4.9	3.2	.....	.....	15	857	857	.....	.....
29	.....	.....	3,282	1,641	17	635	635	.....	.....
30	5.1	5.1	.....	.....	19	543	543	.....	.....
Aug. 1	44	44	4,978	3,318	.....	.....	.....	.....	.....

The period of incubation of the fungus being from 11 to 13 days, as shown by artificial infection experiments, a corresponding increase in the number of spots on the leaves would not necessarily follow immediately after a period during which conditions favorable for infection prevailed.

Notwithstanding the early appearance of the spots in the medium-early field—on June 20—and a consequent expectation of an epidemic of the disease, the increase in infection was very light (Table VIII)

during the latter part of June and early part of July. This was doubtless due to the fact that during this period the stomata were closed against the fungus the greater part of the day on account of the excessively high temperature, which was generally above 100° F., and the excessively low humidity, which at one time fell to 10 and which was only above 60 from 6 to 15 hours a day, and also to the fact that a temperature as high as 95° inhibits the growth of the fungus and kills it after a few days.

After July 5 the temperature was lower and the humidity higher than during the period above mentioned. As a result, numerous conidia were produced from July 9 to 12, and at the end of the period of incubation—July 21 to 25—there was a slight increase in the number of spots (Table VIII). Rains between July 19 and 25 and the resulting high humidity caused a rather marked increase in the number of spots in late July and early August, these spots appearing on many leaves hitherto uninfected (Table VI). Prior to this period the number of leaves showing spots were comparatively few, but after July 30 the majority showed spots, and the proportion of infected to uninfected leaves gradually increased until August 28, after which it decreased. During the period from July 30 to August 28 the humidity was comparatively high, remaining above 60 on an average of 14.6 hours on all except three days, on which it remained above 60 for 10 hours; and the maximum and minimum temperatures generally were not above 90 nor below 55, respectively. The increased proportion of infected to uninfected leaves during this period, however, was not necessarily due to increasingly favorable climatic conditions but to the cumulative effect of the organism, the amount of viable conidia and consequent new infections increasing as the number of spots increased, as shown by the enormous increase of 3,776 spots per plant on August 27 and 28. After September 3 the increase in infection was considerably less (fig. 9). This was due apparently to the fall in temperature, the maximum being rarely above 76° F and the minimum seldom above 50° after September 8, while the humidity was comparatively favorable.

The increase in infection through the season was considerably higher in the early than in the medium-early sugar-beet field, as shown by the total amount of the disease (Tables VI and VII) and the actual increase (Table VIII and fig. 9). This was due to the fact that the foliage was heavier in the early than in the medium-early field (Tables VI and VII, functional leaves), and consequently the humidity was higher and the infection greater in the former than in the latter (Table VIII and fig. 9).

The maximum increase in spots was reached on August 11 in the early field and on August 28 in the medium-early sugar-beet field. The period of greatest increase in the disease is not its period of greatest destructiveness, however, as the plant is not immediately affected by the disease, some time being required for the leaves to be killed.

Prior to August 1, only isolated records of humidity were made in the early field,<sup>1</sup> but after this date continuous records of both humidity and temperatures in both fields were available for comparison (fig. 10).<sup>2</sup> The temperatures prevailing in the two sugar-beet fields were quite comparable, but the humidity was generally different. For instance, from August 2 to 23 the humidity remained above 60 for a longer time, and the maximum humidity was, as a rule, higher in the early than in the medium-early field; from August 23 to September 1 the maximum humidity was lower in the early than in the medium-early field; after the latter date strikingly lower, the difference ranging from 5 to 15 units; after September 5 the humidity remained above 60 for a shorter time in the former than in the latter field; but from September 6 to 21 the range of humidity in the two fields was much closer than during the periods previously mentioned.

The difference in the humidity of the two fields seemed to be due to the difference in the amount of foliage present. Early in the experiment the foliage was heavier in the early than in the medium-early field, but owing to an extremely severe infection, which developed between July 29 and August 13 (fig. 9), the relative proportion of foliage in the two fields was reversed after that period. As a result of this reversal, less moisture was retained and the humidity was lower in the early than in the medium-early field during September, and consequently at that time the relative increase in infection was less in the former than in the latter. Speaking more specifically, early in the experiment there was an average of 29 functional leaves per plant in the early field and 22 in the medium-early; on August 15 there was an average of 26 leaves per plant in both fields, while later on there were fewer per plant in the early than in the medium-early field. On the other hand, on August 13 there was an average of 23.5 infected leaves per plant, with an average of 710 spots per leaf in the early field, and on September 8 there was an average of 21.5 infected leaves per plant, with an average of 508.3 spots per leaf, in the medium-early field.

A comparison of the death rate of the leaves in the two fields before and after the disease appeared shows its destructiveness. For instance, in the early and medium-early fields, from July 7 to 29 and from July 2 to August 25, when no leaves were killed by the fungus, the death rate from normal causes was approximately one leaf per plant in three and four days, respectively; while from July 29 to August 15 and August 25 to September 19, when the disease was most severe in the two fields, the death rate averaged one leaf per plant in nine-tenths of a day and one and three-tenths days, respectively.

<sup>1</sup> These and the later continuous records indicate that prior to August 1 the humidity was generally higher in the early than in the medium-early field.

<sup>2</sup> The temperature records taken at the Weather Bureau station were included in the comparisons and were found to agree closely with those obtained in the two fields.




FIG. 1.—Curves of the maximum and minimum temperature and relative humidities and the number of hours that the humidity remained above 60 from noon of the preceding noon of the given day among the sugar-beet plants of a medium-early and an early field. For comparison, the maximum and minimum temperature records from the Weather Bureau station are included. August 2 to September 2, 1913, at Rocky Ford, Colo.

## SUMMARY

(1) The life of the fungus *Cercospora beticola* overwintering in sugar-beet-top material varies with different environment. When exposed to outdoor conditions, the conidia die in from one to four months; but when kept dry live as long as eight months. The sclerotia-like bodies, which are more or less embedded in the tissues of the host, are more resistant than the conidia, living through the winter when slightly protected, as, for instance, in the interior of a pile of hayed sugar-beet tops or buried in the ground from 1 to 5 inches, and become a source of infection for the succeeding crop. Notwithstanding the difference in temperature and soil-moisture conditions, similar results from the overwintering experiments were obtained at Rocky Ford, Colo., and Madison, Wis.

(2) Climatic conditions and the development of the leafspot can be correlated only when all records are taken at the same relative positions, as shown by comparisons of the Weather Bureau records and the records taken among the plants and 5 feet above the field.

(3) The maximum temperature is much higher near the ground than 5 feet above early in the season, but the difference diminishes as the season advances.

(4) Throughout the season the maximum relative humidity was higher among the leaves than 5 feet above the field. Early in the season, while the plants were small, the humidity remained above 60 longer each day 5 feet above the field than among the plants near the ground; but after the plants attained a good size this condition was reversed. Because of this difference, only records collected among the leaves should be considered in correlating climatic conditions and conidial production and infection.

(5) The effect of rainfall and irrigation on the increase of relative humidity and its duration is apparently much the same.

(6) Thermal tests with artificial cultures showed (a) that exposure to constant temperatures of 35° and 36° C. is fatal to the growth of the fungus; (b) that growth occurred when cultures after exposure for 3 days to either of these temperatures were changed to 30.8°, and also when they were held at either for 8 hours and then at 20° for 16 hours; and (c) that a temperature of 40.5° was fatal in all combinations tested.

(7) Temperature and relative humidity influence the production of conidia and infection in much the same way. A temperature of 80° or 90° F., with a night minimum preferably not below 60°, is most favorable to conidial production, while it is checked by a temperature of 100° or higher and greatly checked by a range from below 50° to 80°. A maximum humidity ranging above 60 for not less than 15 to 18 hours each day induces a good growth of the fungus.

(8) Because of the higher humidity on the lower than on the upper surface of the leaf, the conidia are generally more abundant on the lower surface of the spots, but because of the action of rain and wind they disappear more rapidly from the upper surface.

## LITERATURE CITED

FRÖDIN, John.

1913. Beobachtungen über den Einfluss der Pflanzendecke auf die Bodentemperatur. *In* Lunds Univ. Årsskr., n. f. afd. 2, bd. 8, no. 9, 16 p., 1 fig., 4 pl. Literatur, p. 15.

MASSEE, George.

1906. Plant diseases.—IV. Diseases of beet and mangold. *In* Roy. Gard. Kew, Bul. Misc. Inform., 1906, no. 3, p. 49-60, 5 fig.

PAMMEL, L. H.

1891. Fungus diseases of sugar beet. *In* Iowa Agr. Exp. Sta. Bul. 15, p. 234-254, 7 pl.

POOL, Venus W., and MCKAY, M. B.

1915. Phoma betae on the leaves of the sugar beet. *In* Jour. Agr. Research, v. 4, no. 2, p. 169-178, pl. 27.

1916. Relation of stomatal movement to infection by *Cercospora beticola*. *In* Jour. Agr. Research, v. 5, no. 22, p. 1011-1038, 6 fig., pl. 80-81.

SHAW, H. B.

1914. An improved cog psychrometer. *In* Plant World, v. 7, no. 6, p. 183-185, 2 fig.

STEWART, F. C.

1913. The persistence of the potato late-blight fungus in the soil. N. Y. Agr. Exp. Sta. Bul. 367, p. 357-361.

THÜMMEN, Felix von.

1886. Die Bekämpfung der Pilzkrankheiten unserer Culturgewächse. 137 p. Wien.

TREBOUX, O.

1914. Überwinterung vermittels Mycel bei einigen parasitischen Pilzen. *In* Mycol. Centbl., Bd. 5, Heft 3, p. 120-126.






PLATE III

*Cercospora beticola*: Overwintering tests on the experimental field at Rocky Ford, Colo., during 1912-13:

Sugar-beet leaves infected with *Cercospora beticola* (7) stored in soil in boxes, (2), buried in the ground at different depths from 1 to 8 inches, and (3) left exposed above the ground in a pile of hayed sugar-beet tops.






Fig. 1. Agricultural equipment

Fig. 2. Fig. 3.

#### PLATE IV

Field stations for the collection of weather data at Rocky Ford, Colo., in 1913:

Fig. 1.—Weather shelter, anemometer, and rain gauge at edge of sugar-beet field.

Fig. 2.—Weather shelter among beet plants, showing hygrothermograph and cog psychrometer.

Fig. 3.—Weather shelter of the local Weather Bureau station about 3 miles from sugar-beet field.

ADDITIONAL COPIES  
OF THIS PUBLICATION MAY BE PROCURED FROM  
THE SUPERINTENDENT OF DOCUMENTS  
GOVERNMENT PRINTING OFFICE  
WASHINGTON, D. C.  
AT  
20 CENTS PER COPY  
SUBSCRIPTION PRICE, 3.00 PER YEAR  
V

